

A detailed 3D rendering of a cell membrane. The top portion shows a lipid bilayer with various receptors and signaling molecules. Below the membrane, a large number of blue, spherical cells are arranged in a grid-like pattern. Some of these cells are highlighted with different colors (pink, orange, purple) and contain internal structures, suggesting they are in different states or stages of a process. The overall scene is illuminated with a mix of red, blue, and purple light, creating a vibrant and scientific atmosphere.

biotechne® / R&D SYSTEMS

Products for IL-1 Family Research

The IL-1 Cytokine Family

The IL-1 cytokine family consists of eleven members, including seven cytokines with pro-inflammatory activities (IL-1 α , IL-1 β , IL-18, IL-33, IL-36 α , IL-36 β , and IL-36 γ), one with anti-inflammatory activity (IL-37), and three receptor antagonists (IL-1ra, IL-36Ra, and IL-38). Cytokines belonging to the IL-1 family share a conserved β -trefoil structure and bind to receptors belonging to the IL-1 receptor family (IL-1 R1-IL-1 R10). With the exception of IL-1ra/IL-1F3, IL-1 family cytokines lack a signal peptide and therefore are not thought to be secreted by the conventional endoplasmic reticulum/Golgi-dependent secretory pathway used by other cytokines. IL-1 β and IL-18 are synthesized as inactive precursor proteins that are activated and secreted following cleavage by Caspase-1. N-terminal processing is also required for IL-36 α , IL-36 β , IL-36 γ , and IL-36Ra to be fully active, but the processing enzymes involved have not been identified.

IL-1 family cytokines activate intracellular signaling pathways by binding to a primary receptor subunit, such as IL-1 RI/IL-1 R1, IL-18 R α /IL-1 R5, IL-1 Rrp2/IL-1 R6, or ST2/IL-1 R4, which then recruits a co-receptor to form the active receptor complexes. Most IL-1 family receptors have three extracellular immunoglobulin-like domains and a cytoplasmic Toll/IL-1 receptor (TIR) domain, with the exceptions of SIGIRR/IL-1 R8, which has only one Ig-like domain and IL-1 RII/IL-1 R2, which lacks an intracellular TIR domain. The TIR domain is also conserved in Toll-like receptors (TLRs) and a number of intracellular adaptor proteins that mediate IL-1 R/TLR signaling. One additional structural difference among IL-1 family receptors is that SIGIRR/IL-1 R8, TIGIRR-1/IL-1 R9, and TIGIRR-2/IL-1 R10 contain

an intracellular C-terminal extension that is not found in other receptors belonging to this family. Signaling cascades triggered by the seven pro-inflammatory IL-1 family cytokines activate MAPKs, AP-1, and NF κ B, leading to the expression of pro-inflammatory cytokines, chemokines, and secondary mediators of the inflammatory response. In addition, many of these cytokines have been shown to regulate the differentiation and function of T helper cells. Other members of the IL-1 family inhibit inflammation by functioning as antagonists of IL-1 or IL-36 signaling. IL-1ra negatively regulates IL-1 signaling by binding to IL-1 RI/IL-1R1, which inhibits its ability to interact with IL-1 α and IL-1 β . Similarly, IL-36Ra binds to IL-1 Rrp2/IL-1 R6 and inhibits IL-36 signaling. Both the IL-1 and IL-36 receptor antagonists are incapable of initiating downstream signaling on their own due to an inability to recruit the IL-1 RAcP/IL-1 R3 co-receptor. Although less is currently known about IL-37 and IL-38, both are also thought to have anti-inflammatory effects. Five splice variants of IL-37 exist, with four containing a putative Caspase-1 cleavage site. Both the immature and the mature forms of the longest isoform, IL-37b, bind to IL-18 BP and enhance its ability to inhibit IL-18 activity. Additionally, IL-37 binds to a receptor complex consisting of IL-18 R α /IL-1 R5 and SIGIRR/IL-1 R8 and has anti-inflammatory effects. Like the precursor forms of IL-1 α and IL-33, the mature form of IL-37b can also translocate to the nucleus where it may act as a transcriptional regulator. IL-38 binds to the IL-36 receptor, IL-1 Rrp2/IL-1 R6, and soluble IL-1 RI/IL-1 R1. Initial data suggests that IL-38 has antagonistic effects similar to those induced by IL-36Ra. For additional information, please visit our website at bio-techne.com/il-1family.

IL-1 Family Cytokines

| Cytokine | Alternate Name | Receptor | Co-receptor | Function |
|----------------|----------------|-------------------------------------|--------------------------|---|
| IL-1 α | IL-1F1 | IL-1 RI/IL-1 R1 or IL-1 RII/IL-1 R2 | IL-1 RAcP/IL-1 R3 | Pro-inflammatory |
| IL-1 β | IL-1F2 | IL-1 RI/IL-1 R1 or IL-1 RII/IL-1 R2 | IL-1 RAcP/IL-1 R3 | Pro-inflammatory |
| IL-1ra | IL-1F3 | IL-1 RI/IL-1 R1 | | Receptor antagonist; inhibits inflammation |
| IL-18 | IL-1F4 | IL-18 R α /IL-1 R5 | IL-18 R β /IL-1 R7 | Pro-inflammatory |
| IL-33 | IL-1F11 | ST2/IL-1 R4 | IL-1 RAcP/IL-1 R3 | Pro-inflammatory |
| IL-36 α | IL-1F6 | IL-1 Rrp2/IL-1 R6 | IL-1 RAcP/IL-1 R3 | Pro-inflammatory |
| IL-36 β | IL-1F8 | IL-1 Rrp2/IL-1 R6 | IL-1 RAcP/IL-1 R3 | Pro-inflammatory |
| IL-36 γ | IL-1F9 | IL-1 Rrp2/IL-1 R6 | IL-1 RAcP/IL-1 R3 | Pro-inflammatory |
| IL-36 Ra | IL-1F5 | IL-1 Rrp2/IL-1 R6 | | Receptor antagonist; inhibits inflammation |
| IL-37 | IL-1F7 | IL-18 R α /IL-1 R5 | SIGIRR/IL-1 R8 | Anti-inflammatory |
| IL-38 | IL-1F10 | IL-1 Rrp2/IL-1 R6 | | Putative receptor antagonist; inhibits inflammation |


IL-1 Receptor Family

| Receptor | Co-receptor | Function |
|------------------|-------------|--|
| IL-1 RI | IL-1 R1 | Primary cytokine-binding receptor for IL-1 α and IL-1 β ; inhibited by IL-1ra binding |
| IL-1 RII | IL-1 R2 | Decoy receptor; inhibits IL-1 signaling |
| IL-1 RAcP | IL-1 R3 | Co-receptor for IL-1 RI, ST2/IL-1 R4, and IL-1 Rrp2/IL-1 R6 |
| ST2/IL-33 R | IL-1 R4 | Primary cytokine-binding receptor for IL-33 |
| IL-18 R α | IL-1 R5 | Primary cytokine-binding receptor for IL-18 |
| IL-1 Rrp2 | IL-1 R6 | Primary cytokine-binding receptor for IL-36 α , IL-36 β , and IL-36 γ ; inhibited by IL-36Ra binding |
| IL-18 R β | IL-1 R7 | Co-receptor for IL-18 R α /IL-1 R5 binding to IL-18 |
| SIGIRR/TIR8 | IL-1 R8 | Co-receptor for IL-18R α /IL-1 R5 binding to IL-37 |
| TIGIRR-1 | IL-1 R9 | Orphan receptor |
| TIGIRR-2 | IL-1 R10 | Orphan receptor |

IL-1 Cytokine Family Inhibitors



IL-1 Inhibitors

| | |
|--|---|
| IL-1ra | An IL-1 family cytokine that acts as an IL-1 receptor antagonist by competing with IL-1 α and IL-1 β for binding to IL-1 RI/IL-1 R1; binding of IL-1ra to IL-1 RI/IL-1 R1 inhibits recruitment of IL-1 RAcP and downstream signaling. |
| IL-1 RII/IL-1 R2 | An IL-1 decoy receptor with a short cytoplasmic domain that is incapable of transducing an IL-1 signal; binds to pro-IL-1 α and pro-IL-1 β in the cytosol, preventing their cleavage and activation by different enzymes. |
| SIGIRR/TIR8 | Single immunoglobulin domain-containing IL-1 receptor-related (SIGIRR) molecule; a subtype of the IL-1 receptor family that contains a single extracellular immunoglobulin-like domain and may inhibit signaling by IL-1 family cytokines in a context-dependent manner by preventing association of the co-receptor or interfering with the association of TIR-containing adaptor proteins with the receptor complex; co-receptor for IL-37. |
| Soluble IL-1 RI/IL-1 R1 or IL-1 RII/IL-1 R2 | Soluble receptors that can bind to IL-1 and IL-1 RAcP/IL-1 R3 but are incapable of propagating a signal. |
| Soluble IL-1 RAcP/IL-1 R3 | A soluble receptor that can bind to IL-1 - IL-1 RI/IL-1 R1 but is incapable of propagating a signal; enhances IL-1 binding to soluble IL-1 RII/IL-1 R2. |



IL-18 Inhibitors

| | |
|---|--|
| IL-18 BP | IL-18 binding protein (IL-18 BP); a soluble protein that binds to IL-18 with higher affinity than either the cell-bound or soluble forms of IL-18 R and prevents IL-18 signaling |
| SIGIRR/TIR8 | Single immunoglobulin domain-containing IL-1 receptor-related (SIGIRR) molecule; a subtype of the IL-1 receptor family that contains a single extracellular immunoglobulin-like domain and may inhibit signaling by IL-1 family cytokines in a context-dependent manner by preventing association of the co-receptor or interfering with the association of TIR-containing adaptor proteins with the receptor complex; co-receptor for IL-37 |
| Soluble IL-18 Rα | A soluble receptor that can bind to IL-18 but is incapable of propagating a signal; a weak inhibitor compared to IL-18 BP |



IL-1 Family Signaling Pathways

Click on one of the IL-1 family cytokines below to highlight the signaling pathway and overall effect induced by each cytokine along with the intrinsic inhibitors that may alter its activity.

Interactive Pathways

- IL-1
- IL-1ra
- IL-18
- IL-33
- IL-36
- IL-36Ra
- IL-37
- IL-38

[View Full Pathway](#)

[Request Print Version](#)

IL-1 Inhibitors

IL-1ra
An IL-1 family cytokine that acts as an IL-1 receptor antagonist by preventing IL-1 α or IL-1 β from binding to IL-1 RI. Binding of IL-1ra to IL-1 RI inhibits recruitment of IL-1 RAcP and downstream signaling.

IL-1 RII
An IL-1 decoy receptor with a short cytoplasmic domain that is incapable of transducing an IL-1 signal.

SIGIRR/TIR8
Single immunoglobulin domain-containing IL-1 receptor-related (SIGIRR) molecule. A subtype of the IL-1 R family that contains a single extracellular immunoglobulin-like domain and may inhibit signaling by IL-1 family cytokines in a context-dependent manner by preventing association of the co-receptor or interfering with the association of TIR-containing adaptor proteins with the receptor complex; co-receptor for IL-37.

Soluble IL-1 RAcP
A soluble receptor that can bind to IL-1 - IL-1 RI but is incapable of propagating a signal. Enhances IL-1 binding to soluble IL-1 RII.

Soluble IL-1 RI or RII
Soluble receptors that can bind to IL-1 and IL-1 RAcP but are incapable of propagating a signal.



Interact with this Pathway

Scan the QR Code or visit:
rnds.com/pathways_il-1familysignaling


IL-33 Inhibitors

SIGIRR/TIR8

Single immunoglobulin domain-containing IL-1 receptor-related (SIGIRR) molecule; a subtype of the IL-1 receptor family that contains a single extracellular immunoglobulin-like domain and may inhibit signaling by IL-1 family cytokines in a context-dependent manner by preventing association of the co-receptor or interfering with the association of TIR-containing adaptor proteins with the receptor complex; co-receptor for IL-37

Soluble ST2/IL-1R4

A soluble receptor that can bind to IL-33 but is incapable of propagating a signal




IL-36 Inhibitors

IL-36Ra

An IL-1 family cytokine that acts as an IL-36 receptor antagonist by preventing IL-36 α , IL-36 β , or IL-36 γ from binding to IL-1 Rrp2/IL-1 R6; binding of IL-36ra to IL-1 Rrp2/IL-1 R6 inhibits recruitment of IL-1 RAcP/IL-1 R3 and downstream signaling

SIGIRR/TIR8

Single immunoglobulin domain-containing IL-1 receptor-related (SIGIRR) molecule; a subtype of the IL-1 receptor family that contains a single extracellular immunoglobulin-like domain and may inhibit signaling by IL-1 family cytokines in a context-dependent manner by preventing association of the co-receptor or interfering with the association of TIR-containing adaptor proteins with the receptor complex; co-receptor for IL-37





R&D Systems™ Products for IL-1 Family Research

Recombinant Proteins

R&D Systems has an unparalleled selection of bioactive proteins for IL-1 family research. We offer recombinant human and mouse proteins for most of the IL-1 family cytokines and receptors including bioactive N-terminally truncated forms of Recombinant Human and Mouse IL-36 α , IL-36 β , and IL-36 γ , and Recombinant Human IL-37b. In addition, we offer Recombinant Human and Mouse IL-18. Stringent production and purification standards, along with our rigorous bioassay testing, ensure that R&D Systems proteins will provide industry-leading bioactivity and lot-to-lot consistency.

R&D Systems Offers the Widest Selection of Pro- and Anti-Inflammatory IL-1 Family Cytokines




IL-36 β and IL-36 γ Induce IL-8 Secretion by A431 Cells. The A431 human epithelial carcinoma cell line was treated with increasing concentrations of Recombinant Human IL-36 β /IL-1F8 (aa 5-157; Catalog # 6834-ILB; orange line) or Recombinant Human IL-36 γ /IL-1F9 (aa 18-169; Catalog # 6835-IL; green line). The levels of CXCL8/IL-8 in the cell culture supernatants were measured using the Human CXCL8/IL-8 DuoSet® ELISA Development System (Catalog # DY208). The ED₅₀ for this effect is 0.8-4.8 ng/mL following treatment with Recombinant Human IL-36 β /IL-1F8 and 1.5-9 ng/mL following treatment with Recombinant Human IL-36 γ /IL-1F9.

IL-37b Enhances the Inhibition of IL-18-induced IFN- γ Release by IL-18 Binding Protein. The KG-1 human acute myelogenous leukemia cell line was treated with 40 ng/mL Recombinant Human IL-18 (Catalog # 9124-IL), 5 ng/mL Recombinant Human IL-18 BP α (Catalog # 119-BP), and increasing concentrations of Recombinant Human IL-37b/IL-1F7 β (Catalog # 7585-IL). The levels of IFN- γ in cell culture supernatants were measured using the Human IFN- γ Quantikine® ELISA Kit (Catalog # DIF50C). The ED₅₀ for this effect is 2.5-12.5 μ g/mL.


Recombinant Human and Mouse IL-18

Benefits Include:

- High Activity Levels that are Equivalent to the Leading Competitor's Proteins
- Increased Stability Compared to the Leading Competitor
- Superior Purity



Recombinant Human IL-18 Induces IFN- γ Production by KG-1 Cells. The KG-1 human acute myelogenous leukemia cell line was treated with increasing concentrations of Recombinant Human IL-18/IL-1F4 (Catalog # 9124-IL) and the levels of IFN- γ in cell culture supernatants were assessed using the Human IFN- γ Quantikine[®] ELISA Kit (Catalog # DIF50C). The ED₅₀ for this effect is 1.5–9 ng/mL. The purity of Recombinant Human IL-18/IL-1F4 (Catalog # 9124-IL) was assessed by SDS-PAGE analysis under reducing (R) and non-reducing (NR) conditions and visualized by silver staining.



Recombinant Mouse IL-18 Induces IFN- γ Production by Activated T Cells. Activated mouse T cells were treated with increasing concentrations of Recombinant Mouse IL-18/IL-1F4 (Catalog # 9139-IL) and the levels of IFN- γ in cell culture supernatants were measured using the Mouse IFN- γ Quantikine[®] ELISA Kit (Catalog # MIF00). The ED₅₀ for this effect is 0.06–0.36 ng/mL. The purity of Recombinant Mouse IL-18/IL-1F4 (Catalog # 9139-IL) was assessed by SDS-PAGE analysis under reducing (R) and non-reducing (NR) conditions and visualized by silver staining.

Recombinant Proteins

IL-1 Family Cytokines

| Molecule | Species | Catalog # |
|------------------------------------|---------|-----------|
| IL-1 α /IL-1F1 | Human | 200-LA |
| | Mouse | 400-ML |
| | Rat | 500-RL |
| IL-1 β /IL-1F2 | Human | 201-LB |
| | Mouse | 401-ML |
| | Rat | 501-RL |
| IL-1 τ a/IL-1F3 | Human | 280-RA |
| | Mouse | 480-RM |
| | Rat | 1545-RA |
| IL-18/IL-1F4 | Human | 9124-IL |
| | Mouse | 9139-IL |
| | Rat | 521-RL |
| IL-33/IL-1F11 | Human | 3625-IL |
| | Mouse | 3626-ML |
| IL-36 α /IL-1F6 (aa 1-158) | Human | 1078-IL |
| IL-36 α /IL-1F6 (aa 6-158) | Human | 6995-IL |
| IL-36 α /IL-1F6 (aa 1-160) | Mouse | 2297-ML |
| IL-36 α /IL-1F6 (aa 8-160) | Mouse | 7059-ML |
| IL-36 β /IL-1F8 (aa 1-157) | Human | 1099-IL |
| IL-36 β /IL-1F8 (aa 5-157) | Human | 6834-ILB |
| IL-36 β /IL-1F8 (aa 1-183) | Mouse | 2298-ML |
| IL-36 β /IL-1F8 (aa 31-183) | Mouse | 7060-ML |
| IL-36 γ /IL-1F9 (aa 1-169) | Human | 2320-IL |
| IL-36 γ /IL-1F9 (aa 18-169) | Human | 6835-IL |
| IL-36 γ /IL-1F9 (aa 13-164) | Mouse | 6996-IL |
| IL-36Ra/IL-1F5 | Human | 1275-IL |
| | Mouse | 2714-ML |
| IL-37/IL-1F7 | Human | 1975-IL |
| IL-37b/IL-1F7b | Human | 7585-IL |

IL-1 Family Receptors

| Molecule | Species | Catalog # |
|---------------------------|---------|-----------|
| IL-1 RI/IL-1 R1 | Human | 269-1R |
| | Mouse | 771-MR |
| | Rat | 4895-MR |
| IL-1 RII/IL-1 R2 | Human | 263-2R |
| | Mouse | 663-2R |
| IL-1 RAcP/IL-1 R3 | Human | 9176-CP |
| | Human | 1007-MR |
| IL-1 Rrp2/IL-1 R6 | Human | 872-RP |
| | Mouse | 2354-RP |
| | Rat | 573-RP |
| IL-18 R α /IL-1 R5 | Human | 816-LR |
| IL-18 R β /IL-1 R7 | Human | 118-AP |
| SIGIRR | Human | 990-SG |
| | Mouse | 992-SG |
| ST2/IL-33 R | Human | 523-ST |
| | Mouse | 1004-MR |


Recombinant proteins for additional species are also available for most molecules. Please visit our website at [bio-techne.com/il-1family](https://www.bio-techne.com/il-1family) for a complete product listing.




Learn More

Scan the QR Code or visit:

R&D Systems Also Offers IL-1 Family Receptor Antagonists



Recombinant Human IL-1ra Inhibits IL-1 α -induced Cell Proliferation. The D10.G4.1 mouse helper T cell line was treated with 50 pg/ml Recombinant Human IL-1 α /IL-1F1 (Catalog # 200-LA) and increasing concentrations of Recombinant Human IL-1ra/IL-1F3 (Catalog # 280-RA) and cell proliferation was assessed. The ED₅₀ for this effect is 5–40 ng/mL.




Recombinant Human IL-36Ra Inhibits IL-36 β -induced IL-8 Secretion. The A431 human epithelial carcinoma cell line was treated with 10 ng/mL Recombinant Human IL-36 β (Catalog # 6834-ILB) and increasing concentrations of Recombinant Human IL-36Ra/IL-1F5 (Catalog # 1275-IL). The levels of CXCL8/IL-8 in the cell culture supernatants were measured using the Human CXCL8/IL-8 DuoSet™ ELISA Development System (Catalog # DY208). The ED₅₀ for this effect is 0.2–1 µg/mL.


Unlabeled and Fluorochrome-conjugated Antibodies for IL-1 Family Cytokines and Receptors

R&D Systems offers both unconjugated and fluorochrome-conjugated antibodies for studying IL-1 family cytokines and their receptors. These antibodies are qualified for blocking/neutralization, flow cytometry, immunocytochemistry/immunohistochemistry, and/or Western blot. All of our antibodies are 100% guaranteed to work in the application and species listed on the R&D Systems website.


Antibodies for Blocking/Neutralization



IL-1 α -induced Cell Proliferation and Neutralization using an Anti-Human IL-1 α Antibody. Proliferation of the D10.G4.1 mouse helper T cell line was assessed following treatment with increasing concentrations of Recombinant Human IL-1 α /IL-1F1 (Catalog # 200-LA; orange line). Proliferation stimulated by 50 pg/mL Recombinant Human IL-1 α /IL-1F1 was neutralized by treating the cells with increasing concentrations of a Goat Anti-Human IL-1 α /IL-1F1 Antigen Affinity-purified Polyclonal Antibody (Catalog # AF-200-NA; green line). The ND₅₀ for this effect is typically 4–20 ng/mL in the presence of 1.25 µg/mL concanavalin A.



IL-33-induced Cell Proliferation and Neutralization using an Anti-Human IL-33 Antibody. Proliferation of the D10.G4.1 mouse helper T cell line was assessed following treatment with increasing concentrations of Recombinant Mouse IL-33 (Catalog # 3626-ML; orange line). Proliferation stimulated by 0.25 ng/mL Recombinant Mouse IL-33 was neutralized by treating the cells with increasing concentrations of a Goat Anti-Mouse IL-33 Antigen Affinity-purified Polyclonal Antibody (Catalog # AF3626; green line). The ND₅₀ for this effect is typically 10–50 ng/mL.



IL-33-induced IFN- γ Secretion and Neutralization using an Anti-Human ST2/IL-33R Antibody. Human peripheral blood mononuclear cells were treated with 0.25 ng/mL Recombinant Human IL-12 (Catalog # 219-IL) and increasing concentrations of Recombinant Human IL-33 (Catalog # 3625-IL). IFN- γ secretion was measured using the Human IFN- γ Quantikine™ ELISA Kit (Catalog # DIF50C; orange line). IFN- γ secretion induced by 1 ng/mL Recombinant Human IL-33 was neutralized by treating the cells with increasing concentrations of a Goat Anti-Human ST2/IL-33 R Antigen Affinity-purified Polyclonal Antibody (Catalog # AF523; green line). The ND₅₀ for this effect is typically 0.1–0.6 µg/mL.




Learn More


Scan the QR Code or visit:

bio-techne.com/il-1family


Antibodies for Immunocytochemistry or Flow Cytometry



Detection of ST2/IL-1 R4 in KG-1 Cells. The KG-1 human acute myelogenous leukemia cell line was stained with an APC-conjugated Goat Anti-Human ST2/IL-1 R4 Antigen Affinity-purified Polyclonal Antibody (Catalog # FAB5231A; filled histogram) or an APC-conjugated Goat IgG Isotype Control (Catalog # IC108A; open histogram).



Detection of IL-1β/IL-1F2 in MCF-7 Cells. IL-1β/IL-1F2 was detected in the immersion-fixed MCF-7 human breast cancer cell line using a Goat Anti-Mouse IL-1β/IL-1F2 Antigen Affinity-purified Polyclonal Antibody (Catalog # AF-401-NA) at 8 μg/mL for 3 hours at room temperature. Cells were stained using the NorthernLights™ 557-conjugated Anti-Goat IgG Secondary Antibody (Catalog # NL001; red) and counterstained with DAPI (blue). Specific staining was localized to the cytoplasm.



Detection of IL-18 Rα/IL-1 R5 in Human Peripheral Blood Mononuclear Cells by Flow Cytometry. Human peripheral blood mononuclear cells were treated with 5 μg/mL PHA and 10 ng/mL Recombinant Human IL-2 for 2 days and then stained with an APC-conjugated Mouse Anti-Human IL-18 Rα/IL-1 R5 Monoclonal Antibody (Catalog # FAB840A; filled histogram) or an APC-conjugated Mouse IgG, Isotype Control Antibody (Catalog # IC002A; open histogram).

Select R&D Systems Antibodies for Studying IL-1 Family Cytokines & Their Receptors

IL-1 Family Cytokines

| Molecule | Species | Clone | Unlabeled Antibodies Catalog # (Applications) |
|----------------------------|------------|--------------------------------|---|
| IL-1α/IL-1F1 | Human | 4414 | MAB200 (B/N, E, ICC/IF, WB) |
| | Human | 4414R | MAB200R (B/N, E, ICC/IF) * |
| | Human | Polyclonal | AF-200-NA (B/N, ICC/IF, WB) |
| | Mouse | 40508 | MAB400 (B/N, E) |
| | Mouse | ALF161 | MAB4001 (B/N, IP, WB) |
| | Mouse | Polyclonal | AF-400-NA (B/N, IHC, WB) |
| | Rat | 59015 | MAB500 (E, WB) |
| Rat | Polyclonal | AF500 (B/N, WB) | |
| IL-1α/IL-1F1 Membrane Form | Human | 3405 | MAB2001 (B/N, FC) |
| IL-1α/IL-1F1 Propeptide | Human | Polyclonal | AF4154 (FC, WB) |
| IL-1β/IL-1F2 | Human | 1027B | MAB8406 (FC) |
| | Human | 8516 | MAB201 (B/N, FC, ICC/IF, WB) |
| | Human | 2805 | MAB601 (B/N, E, ICC/IF, WB) |
| | Human | Polyclonal | AF-201-NA (B/N, ICC/IF, WB) |
| | Mouse | B122 | MAB4012 (B/N, IP, WB) |
| | Mouse | 30311 | MAB401 (B/N, E) |
| | Mouse | 30311R | MAB401R (B/N, ICC/IF) * |
| | Mouse | Polyclonal | AF-401-NA (B/N, ICC/IF, IHC, SW, WB) |
| | Rat | 38123 | MAB5011 (WB) |
| Rat | 38139 | MAB501 (B/N, ICC/IF) | |
| Rat | Polyclonal | AF-501-NA (B/N, E, ICC/IF, WB) | |
| IL-1β/IL-1F2 Propeptide | Human | 615417 | MAB6964 (ICC/IF, SW, WB) |
| IL-1ra/IL-1F3 | Human | 10309 | MAB280 (B/N, E) |
| | Human | Polyclonal | AF-280-NA (B/N, IHC, WB) |
| | Mouse | 694204 | MAB4801 (WB) |
| | Mouse | Polyclonal | AF-480-NA (B/N, E, WB) |
| IL-18/IL-1F4 | Human | 1072F | MAB9124 (B/N) * |
| | Human | 914205 | MAB91241 (B/N) |
| | Human | Polyclonal | AF2548 (B/N, ICC/IF, WB) |
| | Human | 125-2H | D044-3 (B/N, E, IP) |
| | Human | 25-2G | D043-3 (WB) |
| | Mouse | 93-10C | D048-3 (B/N, IP) |
| | Mouse | 39-3F | D046-3 (WB) |
| | Mouse | 74 | D047-3 (E, IP) |
| | Rat | 69604 | MAB521 (WB) |
| | Rat | Polyclonal | AF521 (B/N, WB) |



Learn More

Scan the QR Code or visit:
[bio-technne.com/il-1family](https://www.bio-technne.com/il-1family)

IL-1 Family Cytokines

| Molecule | Species | Clone | Unlabeled Antibodies Catalog # (Applications) |
|---|---------|------------|---|
| IL-18/IL-1F4 Propeptide | Human | 74801 | MAB646 (FC, WB) |
| | Human | Polyclonal | AF646 (WB) |
| IL-33/IL-1F11 | Human | 1061A | MAB36252 (IHC) * |
| | Human | 40015C | MAB36253 (E, IHC, WB) * |
| | Human | 40115D | MAB36254 (B/N, IHC, WB) * |
| | Human | 390412 | MAB3625 (ICC/IF, WB) |
| | Human | Polyclonal | AF3625 (B/N, E, ICC/IF, IHC, WB) |
| | Mouse | 396118 | MAB3626 (FC, ICC/IF, WB) |
| IL-33/IL-1F11 Propeptide | Human | Polyclonal | AF4810 (ICC/IF, WB) |
| | Mouse | 518017 | MAB5010 (WB) |
| | Mouse | Polyclonal | AF5010 (ICC/IF, WB) |
| IL-36α/IL--1F6 | Human | 162122 | MAB1078 (WB) |
| | Human | Polyclonal | AF1078 (IHC, WB) |
| | Mouse | 275339 | MAB2297 (B/N, WB) |
| | Mouse | Polyclonal | AF2297 (B/N, WB) |
| IL-36β/IL-1F8 | Human | 162601 | MAB1099 (WB) |
| | Human | Polyclonal | AF1099 (B/N, IHC, WB) |
| | Mouse | Polyclonal | AF2298 (B/N, WB) |
| IL-36γ/IL-1F9 | Human | 278706 | MAB2320 (B/N, WB) |
| | Human | Polyclonal | AF2320 (B/N, WB) |
| IL-36Ra/IL-1F5 | Human | 190524 | MAB1275 (WB) |
| | Human | Polyclonal | AF1275 (B/N, WB) |
| | Mouse | 759207 | MAB2714 (WB) |
| IL-37/IL-1F7 | Human | 899826 | MAB19751 (FC) |
| | Human | 261506 | MAB1975 (B/N) |
| | Human | Polyclonal | AF1975 (WB) |
| IL-38/IL-1F10 | Human | 316709 | MAB2427 (WB) |
| | Human | Polyclonal | AF2427 (WB) |
| | Mouse | 798036 | MAB7774 (FC, WB) |
| | Mouse | 798036R | MAB7774R (FC, WB) * |

* Indicates a recombinant monoclonal antibody
Application key: B/N Blocking/Neutralization E ELISA FC Flow Cytometry ICC/IF Immunocytochemistry/Immunofluorescence IHC Immunohistochemistry IP Immunoprecipitation SW Simple Western WB Western blot

Antibodies for additional species are available for most target analytes. Please visit our website at bio-techne.com/il-1family for a complete product listing.

IL-1 Family Receptors

| Molecule | Species | Clone | Unlabeled Antibodies Catalog # (Applications) | Fluorochrome-conjugated Antibodies Catalog # (Applications) |
|--|---------|------------|---|---|
| IL-1 RI/IL-1R1 | Human | 35730 | MAB269 (E, WB) | |
| | Human | Polyclonal | AF269 (B/N, FC, WB) | FAB269A, F, N, P (FC) |
| | Mouse | JAMA147 | MAB7711 (B/N) | |
| | Mouse | 129304 | MAB7712 (FC, WB) | FAAB7712F, P (FC) |
| | Mouse | Polyclonal | AF771 (B/N, IHC, WB) | |
| IL-1 RII/IL-1R2 | Human | 32437 | MAB263 (B/N, ICC/IF, WB) | |
| | Human | 34141 | MAB663 (B/N, E, WB) | FAB663A, F, N, P (FC) |
| | Human | Polyclonal | AF-263-NA (B/N, ICC/IF, WB) | |
| | Mouse | 130817 | MAB563 (WB) | |
| IL-1 RAcP/IL-1R3 | Mouse | Polyclonal | AF563 (FC, IHC, WB) | |
| | Human | 89412 | MAB676 (FC) | FAB676A, G, N, P (FC) |
| | Human | 89412R | MAB676R (FC) * | |
| IL-1 RAPL2/IL-1R9 | Human | Polyclonal | AF676 (FC, WB) | |
| | Human | Polyclonal | AF1007 (FC, IHC, WB) | |
| IL-1 Rrp2/IL-1R6 | Mouse | 320017 | MAB3068 (WB) | |
| | Human | Polyclonal | AF872 (WB) | |
| | Mouse | Polyclonal | AF2354 (WB) | |
| | Rat | 131011 | MAB573 (WB) | |
| IL-18 Rα/IL-1R5 | Rat | Polyclonal | AF573 (WB) | |
| | Human | 70614 | MAB8401 (WB) | |
| | Human | 70625 | MAB840 (B/N, FC, ICC/IF, IHC) | FAB840A, G, P (FC) |
| | Human | Polyclonal | AF840 (B/N, FC, IHC, WB) | |
| | Mouse | 112614 | MAB1216 (FC, WB) | FAB1216A, F, N (FC) |
| | Mouse | 112624 | MAB12161 (B/N, WB) | |
| IL-18 Rβ/IL-1R7 | Mouse | Polyclonal | AF856 (B/N, FC, WB) | |
| | Human | 132016 | MAB1181 (B/N, WB) | |
| | Human | 132029 | MAB118 (FC, WB) | FAB118F, P (FC) |
| | Human | Polyclonal | AF118 (B/N, WB) | |
| | Mouse | Polyclonal | AF199 (WB) | |
| SIGIRR/TIR8/IL-1R8 | Human | 162201 | MAB990 (WB) | FAB990A (FC) |
| | Human | Polyclonal | AF990 (WB) | |
| | Mouse | 161917 | MAB1092 (WB) | |
| ST2/IL-33 R | Mouse | Polyclonal | AF1092 (WB) | |
| | Human | 97203 | MAB523 (B/N, E, WB) | |
| | Human | Polyclonal | AF523 (B/N, FC, WB) | FAB5231A, P (FC) |
| | Mouse | 245707 | MAB10041 (B/N, E, FC) | FAB10041A, N, P (FC) |
| | Mouse | 245714 | MAB1004 (WB) | |
| | Mouse | Polyclonal | AF1004 (B/N, WB) | |

* Indicates a recombinant monoclonal antibody
Application key: B/N Blocking/Neutralization E ELISA FC Flow Cytometry ICC/IF Immunocytochemistry/Immunofluorescence IHC Immunohistochemistry IP Immunoprecipitation SW Simple Western WB Western blot
Fluorochrome key for FAB/IC catalog numbers ending in: A Allophycocyanin C PerCP F Fluorescein G AlexaFluor[®] 488 N AlexaFluor[®] 700 P Phycoerythrin
 Alexa Fluor[®] is a registered trademark of Molecular Probes, Inc., Eugene, OR.


ELISA Kits for Detecting IL-1 Family Cytokines and Soluble IL-1 Family Receptors

R&D Systems develops and manufactures the most highly referenced ELISA kits in the world. We offer both complete, ready-to-run Quantikine™ Colorimetric Sandwich ELISA Kits and the more flexible DuoSet™ ELISA Development Systems for detecting IL-1 family cytokines or soluble IL-1 family receptors. Quantikine Kits are rigorously tested in-house to ensure that they provide the highest levels of specificity, accuracy, precision, and sensitivity in analyte quantification without the need for further assay optimization. DuoSet ELISA Development Systems offer an economical alternative to Quantikine Kits by providing all of the components necessary for a customer to develop their own working assay. In addition, we also offer DuoSet IC ELISA Development Systems for quantifying specific intracellular proteins that are activated downstream of IL-1 family cytokines (refer to the product tables on pages 20-25 of this brochure).




Learn More


Scan the QR Code or visit:
bio-techne.com/elisa



Measurement of IL-1 α and IL-1 β in Peripheral Blood Mononuclear Cell Culture Supernatants. Human peripheral blood mononuclear cells were treated with PHA. The levels of IL-1 α /IL-1F1 and IL-1 β /IL-1F2 were assessed using the Human IL-1 α /IL-1F1 Quantikine™ ELISA Kit (Catalog # DLA50) or the Human IL-1 β /IL-1F2 QuantiGlo™ ELISA Kit (Catalog # QLB00B).



Measurement of ST2/IL-1 R4 Levels in Cell Culture Supernatants using the Human ST2/IL-33 R Quantikine® ELISA Kit. Aliquots of cell culture supernatants removed from the BJAB human Burkitt's lymphoma cell line, BUD-8 human skin fibroblast cell line, U87-MG glioblastoma/astrocytoma cell line, HUT-78 human mature cutaneous T cell lymphoma cell line, and human umbilical vein endothelial cells (HUVEC) were assayed for ST2/IL-1 R4 using the Human ST2/IL-33 R Quantikine™ ELISA Kit (Catalog # DST200).



Detection of IL-1 β -induced p38 α Phosphorylation in HepG2 Cells. The HepG2 human hepatocellular carcinoma cell line was treated with Recombinant Human IL-1 β /IL-1F2 (Catalog # 201-LB) for the indicated times. p38 α phosphorylation was assessed in cell lysates using the Human/Mouse/Rat Phospho-p38 α (T180/Y182) DuoSet™ IC ELISA Development System (Catalog # DYC869B; bar graph). The results obtained from the DuoSet IC ELISA are consistent with the relative levels of phosphorylated p38 detected in the same lysates by Western blot (inset).

IL-1 Family Cytokines

| Molecule | Species | Quantikine™ ELISA Kit (Catalog #) | DuoSet™ ELISA Development System or Other ELISA Kit (Catalog #) |
|---------------------------------|-----------|-----------------------------------|---|
| IL-1 α /IL-1F1 | Human | DLA50 | DY200 |
| | Mouse | MLA00 | DY400 |
| | Rat | RRA00 | |
| IL-1 β /IL-1F2 | Human | DLB50* | DY201 |
| | Mouse | MLB00C | DY401 |
| | Rat | RLB00 | DY501 |
| IL-1ra/IL-1F3 | Human | DRA00B | DY280 |
| | Mouse | MRA00 | DY480 |
| IL-18/IL-1F4 | Human | DL180 | DY318 |
| | Mouse | | 7625 |
| IL-18/IL-18 BP α Complex | Human | | DY8936 |
| IL-33/IL-1F11 | Human | D3300B | DY3625B |
| | Mouse/Rat | M3300 | DY3626 |
| IL-36 β /IL-1F8 | Human | | DY1099 |
| IL-36Ra/IL-1F5 | Human | | DY1275 |
| IL-37/IL-1F7 | Human | | DY1975 |
| IL-38/IL-1F10 | Human | | DY9110 |
| | Mouse | | DY2427 |

*Quantikine™ High Sensitivity and QuantiGlo™ ELISA Kits are also available.

IL-1 Family Receptors

| Molecule | Species | Quantikine™ ELISA Kit (Catalog #) | DuoSet™ ELISA Development System (Catalog #) |
|-------------------|---------|-----------------------------------|--|
| IL-1 RI/IL-1 R1 | Human | | DY269 |
| | Mouse | | DY771 |
| IL-1 RII/IL-1 R2 | Human | DR1B00 | DY263 |
| IL-1 RAcP/IL-1 R3 | Human | | DY676 |
| ST2/IL-33 R | Human | DST200 | DY523B |
| | Mouse | MST200 | DY1004 |


Multiplex Assays

In addition to our single analyte ELISA Kits, R&D Systems also offers multiplex assay options for simultaneously detecting multiple target analytes in qualified sample types. These assays include the membrane-based Proteome Profiler™ Antibody Arrays and the bead-based Luminex® Assays and Luminex® High Performance Assays. Several multianalyte profiling kits are available that allow the simultaneous detection of IL-1 α /IL-1F1, IL-1 β /IL-1F2, IL-1ra/IL-1F3, IL-1RI/IL-1R1, and/or IL-1RII/IL-1R2 or intracellular kinases involved in IL-1 family signaling such as MKKs, JNK, and p38.

Proteome Profiler™ Antibody Arrays

Proteome Profiler Antibody Arrays allow for the measurement of up to 119 proteins in a single sample. They require no specialized equipment and eliminate the need for multiple Western blot experiments.

Please visit our website at bio-techne.com/reagents/proteome-profiler-antibody-arrays for more information.




Simultaneous Detection of Multiple Analytes in Lipopolysaccharide-, Nigericin-treated THP-1 Cell Lysates using the Proteome Profiler™ Human Cytokine Array Kit. The THP-1 human acute monocytic leukemia cell line was either untreated or treated with 1 μ g/mL lipopolysaccharide (LPS) for 4 hours and then 5 mM Nigericin (Catalog # 4312) for 1 hour. Cytokine expression in 500 μ L of cell culture supernatant from the untreated and treated cells was analyzed using the Proteome Profiler™ Human Cytokine Array Kit (Catalog # ARY005B).

Luminex® Assays

R&D Systems offers two versions of our Luminex® bead-based assays. Our standard Luminex Assays provide the largest, customizable menu of analytes for bead-based multianalyte profiling using cell culture supernatants, serum, or plasma samples. These assays allow up to 100 analytes to be simultaneously profiled using polystyrene microparticles or 50 analytes using magnetic particles. Please visit bio-techne.com/reagents/luminex-assays/luminex-discovery-assay to see our industry-leading selection of analyte combinations.

Luminex High Performance Assays offer defined analyte panels for bead-based multianalyte profiling. These assays rely on panel-optimized diluents that provide maximum performance for a smaller group of analytes than our standard Luminex Assays. Each assay is fully validated for all sample types indicated for a given panel. In-house testing demonstrates that analyte concentrations determined using our Luminex High Performance Assays correlate closely with those obtained using our single analyte Quantikine™ ELISA Kits. Please visit bio-techne.com/reagents/luminex-assays/luminex-high-performance-assays to view a list of available kits.

Luminex® is a registered trademark of Luminex Corporation



Learn More

Scan the QR Code or visit:
[bio-techne.com/
il-1family](http://bio-techne.com/il-1family)

IL-1 Family Cytokine Signaling

The intracellular signaling pathways triggered by IL-1 α /IL-1F1, IL-1 β /IL-1F2, IL-18/IL-1 F4, IL-33/IL-1 F11, IL-36 α /IL-1F6, IL-36 β /IL-1F8, and IL-36 γ /IL-1F9 activate MAPKs, NF κ B, and AP-1, leading to the expression of pro-inflammatory cytokines, chemokines, and secondary mediators of the inflammatory response. R&D Systems offers a wide selection of antibodies, ELISA Kits, and multiplex assays for studying the intracellular signaling pathways activated downstream of the IL-1 family cytokines and their biological effects.

Select Antibodies, ELISA Kits, and Multiplex Assays for Detecting Signaling Molecules Involved in IL-1 Family Cytokine Signaling

| Molecule | R&D Systems™ or Novus Biologicals™ Antibodies | | R&D Systems™ ELISA/ Assay Kits Catalog # | R&D Systems™ Proteome Profiler™ Antibody Array Kits Catalog # | Tocris™ Small Molecule Activators/ Inhibitors |
|---|---|------------------------------|--|---|---|
| | Species | Catalog # (Applications) | | | |
| ◆ ERK1 | Human | MAB1940 (IHC, WB) | DYC1940 | | ✓ |
| | Human | AF1940 (WB) | | | |
| | Human/ Mouse/Rat | AF1575 (IHC, WB) | | | |
| ◆ Phospho-ERK1 (T202/Y204) | Human/ Mouse/Rat | | DYC1825 | | |
| ◆ ERK1/2 | Human/ Mouse/Rat | MAB1576 (IHC, WB) | | | ✓ |
| | Human/ Mouse/Rat | MAB15761 (WB) | | | |
| | Human/ Mouse/Rat | AF1576 (SW, WB) | | | |
| ◆ Phospho-ERK1 (T202/Y204)/ERK2 (T185/Y187) | Human | MAB1825 (WB) | DYC1018B | ARY003C | |
| | Human/ Mouse | MAB18251 (SW, WB) * | | | |
| | Human/ Mouse/Rat | MAB1018 (FC, ICC/IF, SW, WB) | | | |
| | Human/ Mouse/Rat | IC7806G (FC) | | | |
| | Human/ Mouse/Rat | AF1018 (FC, IHC, SW, WB) | | | |
| Human/ Mouse/Rat | IC1018N (FC) | | | | |

* Indicates a recombinant monoclonal antibody
Application key: B/N Blocking/Neutralization **ChIP** Chromatin Immunoprecipitation
FA Functional Assay **FC** Flow Cytometry **ICC/IF** Immunocytochemistry Immunofluorescence
IHC Immunohistochemistry **IP** Immunoprecipitation **SW** Simple Western **WB** Western blot
Fluorochrome key for FAB/IC catalog numbers ending in: A Allophycocyanin C PerCP
F Fluorescein **G** AlexaFluor® 488 **N** AlexaFluor® 700 **P** Phycoerythrin
◆ Indicates a R&D Systems brand antibody ◆ Indicates a Novus Biologicals brand antibody

| Molecule | R&D Systems™ or Novus Biologicals™ Antibodies | | R&D Systems™ ELISA/ Assay Kits Catalog # | R&D Systems™ Proteome Profiler™ Antibody Array Kits Catalog # | Tocris™ Small Molecule Activators/ Inhibitors |
|--|---|---|--|---|---|
| | Species | Catalog # (Applications) | | | |
| ◆ ERK2 | Human/ Mouse/Rat | MAB1230 (IHC, SW, WB) | DYC1230C | | ✓ |
| | Human/ Mouse/Rat | AF1230 (IHC, WB) | | | |
| ◆ c-Fos | Human | AF7254 (WB) | | | |
| ◆ FosB/GOS3 | Human | AF2214 (IHC, WB) | | | |
| | Human/ Mouse | MAB2214 (WB) | | | |
| ◆ FRA-1 | Human | AF4935 (IHC, WB) | | | |
| | Human | MAB4935 (WB) | | | |
| ◆ I κ B- α | Human | MAB4299 (SW, WB) | DYC4299 | ARY029; ARY027 | ✓ |
| | Human/ Mouse | AF4299 (WB) | | | |
| | Human/ Mouse/Rat | NB100-56507 (FC, ICC/IF, IHC, IP, SW, WB) | | | |
| ◆ Phospho-I κ B- α (S32/S36) | Human | AF4809 (WB) | | | |
| ◆ I κ B- β | Human/Rat | MAB3425 (WB) | | | |
| | Human/ Mouse | AF5225 (WB) | | | |
| ◆ I κ B-e | Human | MAB4300 (WB) | | ARY029; ARY027 | |
| | Human | AF4300 (IHC) | | | |
| | Mouse | AF4637 (WB) | | | |

* Indicates a recombinant monoclonal antibody
Application key: B/N Blocking/Neutralization **ChIP** Chromatin Immunoprecipitation
FA Functional Assay **FC** Flow Cytometry **ICC/IF** Immunocytochemistry Immunofluorescence
IHC Immunohistochemistry **IP** Immunoprecipitation **SW** Simple Western **WB** Western blot
Fluorochrome key for FAB/IC catalog numbers ending in: A Allophycocyanin C PerCP
F Fluorescein **G** AlexaFluor® 488 **N** AlexaFluor® 700 **P** Phycoerythrin
◆ Indicates a R&D Systems brand antibody ◆ Indicates a Novus Biologicals brand antibody



Learn More

Scan the QR Code or visit:
bio-techne.com/il-1family

Select Antibodies, ELISA Kits, and Multiplex Assays for Detecting Signaling Molecules Involved in IL-1 Family Cytokine Signaling

| Molecule | R&D Systems™ or Novus Biologicals™ Antibodies | | R&D Systems™ ELISA/Activity Assay Kits Catalog # | R&D Systems™ Proteome Profiler™ Antibody Array Kits Catalog # | Tocris™ Small Molecule Activators/Inhibitors |
|-----------------------------|---|---|--|---|--|
| | Species | Catalog # (Applications) | | | |
| ◆ IKK-α | Human/Mouse/Rat | AF3768 (ICC/IF, WB) | | ARY029 | ✓ |
| | Human/Mouse | NB100-56704 (FC, ICC/IF, IHC, IP, SW, WB) | | | |
| | Human/Mouse | NBP2-27409 (FC, ICC/IF, IHC, IP, WB) | | | |
| ◆ Phospho-IKK-α (S176/S180) | Human | MAB3768 (WB) | | | |
| ◆ IKK-β | Human | AF4535 (WB) | | ARY029 | ✓ |
| | Human/Mouse | NB100-56509 (FC, ICC/IF, IHC, IP, SW, WB) | | | |
| | Mouse | MAB7155 (WB) | | | |
| ◆ IKK-γ | Human/Mouse/Rat | AF2684 (ICC/IF, SW, WB) | | | ✓ |
| | Human/Mouse/Rat | AF4365 (WB) | | | |
| | Human | NB100-56532 (FC, WB) | | | |
| ◆ IKK-ε | Human | AF3199 (ICC/IF, WB) | | | ✓ |
| | Human/Mouse/Rat | MAB3199 (ICC/IF, WB) | | | |
| ◆ IL-18 BPα | Human | MAB1191 (B/N) | DBP180; DY119 | | |
| | Human | MAB119 (WB) | | | |
| | Human | AF119 (B/N, WB) | | | |
| ◆ IL-18 BPC | Mouse | AF129 (WB) | | | |
| ◆ IL-18 BPD | Mouse | AF122 (B/N, WB) | DY122 | | |
| ◆ IRAK1 | Human | AF4048 (WB) | | ARY029; ARY027 | ✓ |
| ◆ IRAK2 | Human | MAB6690 (WB) | | | |
| ◆ IRAK3 | Human | AF6264 (WB) | | | |
| ◆ IRAK4 | Human | AF3919 (ICC/IF, WB) | | | ✓ |
| ◆ JNK Pan Specific | Human/Mouse/Rat | AF1387 (IHC, WB) | | | ✓ |
| | Human/Mouse/Rat | MAB1387 (WB) | | | |
| ◆ Phospho-JNK Pan Specific | Human/Mouse/Rat | | DYC1387B | ARY003C | |

* Indicates a recombinant monoclonal antibody
Application key: B/N Blocking/Neutralization ChIP Chromatin Immunoprecipitation FA Functional Assay FC Flow Cytometry ICC/IF Immunocytochemistry/Immunofluorescence IHC Immunohistochemistry IP Immunoprecipitation SW Simple Western WB Western blot
Fluorochrome key for FAB/IC catalog numbers ending in: A Allophycocyanin C PerCP F Fluorescein G AlexaFluor® 488 N AlexaFluor® 700 P Phycoerythrin
◆ Indicates a R&D Systems brand antibody ◆ Indicates a Novus Biologicals brand antibody

| Molecule | R&D Systems™ or Novus Biologicals™ Antibodies | | R&D Systems™ ELISA/Activity Assay Kits Catalog # | R&D Systems™ Proteome Profiler™ Antibody Array Kits Catalog # | Tocris™ Small Molecule Activators/Inhibitors |
|----------------------------|---|--------------------------|--|---|--|
| | Species | Catalog # (Applications) | | | |
| ◆ Phospho-JNK (T183/Y185) | Human/Mouse/Rat | AF1205 (IHC, SW, WB) | | ARY018 | |
| | Human/Mouse/Rat | MAB1205 (ICC/IF, SW, WB) | | | |
| ◆ JNK1 | Human/Mouse/Rat | MAB17761 (ICC/IF, WB) | | | ✓ |
| | Human/Mouse/Rat | MAB1776 (WB) | | | |
| ◆ Phospho-JNK1 (T183/Y185) | Human | | | ARY002B | |
| ◆ JNK1/2 | Human/Mouse/Rat | MAB2076 (ICC/IF, WB) | | ARY029 | |
| ◆ JNK2 | Human/Mouse/Rat | MAB1846 (ICC/IF, WB) | | ARY029 | ✓ |
| | Human/Mouse/Rat | AF1846 (WB) | | | |
| ◆ Phospho-JNK2 (T183/Y185) | Human/Mouse/Rat | | DYC2236 | | |
| ◆ c-Jun | Human | MAB2670 (ICC/IF, WB) | | | ✓ |
| | Human | AF2670 (WB) | | | |
| ◆ Phospho-c-Jun (S63) | Human | MAB8930 (ICC/IF, SW, WB) | | ARY003C | |
| ◆ JunB | Human | MAB4456 (WB) | | | |
| | Human | AF4456 (WB) | | | |
| ◆ JunD | Human/Mouse | MAB5526 (WB) | | | |
| | Human/Mouse | AF5526 (WB) | | | |
| ◆ MKK3 | Human/Mouse/Rat | MAB2515 (ICC/IF, WB) | | | |
| ◆ MKK3/MKK6 | Human/Mouse/Rat | MAB2514 (WB) | | | |
| ◆ MKK4 | Human | MAB3390 (ICC/IF) | | | |
| ◆ Phospho-MKK4 (S257/T261) | Human/Mouse/Rat | AF2990 (ICC/IF, WB) | | | |
| ◆ MKK6 | Human/Mouse/Rat | AF16041 (WB) | | | |
| | Human/Mouse/Rat | AF1604 (WB) | | | |

* Indicates a recombinant monoclonal antibody
Application key: B/N Blocking/Neutralization ChIP Chromatin Immunoprecipitation FA Functional Assay FC Flow Cytometry ICC/IF Immunocytochemistry/Immunofluorescence IHC Immunohistochemistry IP Immunoprecipitation SW Simple Western WB Western blot
Fluorochrome key for FAB/IC catalog numbers ending in: A Allophycocyanin C PerCP F Fluorescein G AlexaFluor® 488 N AlexaFluor® 700 P Phycoerythrin
◆ Indicates a R&D Systems brand antibody ◆ Indicates a Novus Biologicals brand antibody

| Molecule | R&D Systems™ or Novus Biologicals™ Antibodies | | R&D Systems™ ELISA/Activity Assay Kits Catalog # | R&D Systems™ Proteome Profiler™ Antibody Array Kits Catalog # | Tocris™ Small Molecule Activators/Inhibitors |
|----------------------------|---|-------------------------------|--|---|--|
| | Species | Catalog # (Applications) | | | |
| ◆ MKK7 | Human | AF3579 (IHC, WB) | | | |
| | Human | MAB3579 (IHC) | | | |
| ◆ MyD88 | Human | AF2928 (FC, ICC/IF, SW, WB) | DY2928 | ARY029 | |
| | Human | MAB2928 (ICC/IF) | | | |
| | Human | MAB29281 (FC) | | | |
| | Human/Mouse | NBP2-27369 (FC, ICC/IF, WB) | | | |
| | Mouse/Rat | AF3109 (FC, ICC/IF, SW, WB) | | | |
| | Mouse | MAB3109 (ICC/IF) | | | |
| ◆ NFκB1 | Human | MAB2697 (WB) | | ARY018; ARY029 | ✓ |
| | Human/Mouse | AF2697 (ChIP, WB) | | | |
| ◆ NFκB2 | Human | MAB28881 (ChIP, ICC/IF, WB) | | ARY029 | ✓ |
| ◆ p38α | Human/Mouse/Rat | AF8691 (IHC, SW, WB) | DYC8691B | | ✓ |
| | Human/Mouse/Rat | MAB869 (WB) | | | |
| ◆ Phospho-p38α (T180/Y182) | Human | MAB8691 (WB) | DYC869B | ARY003C; ARY018 | |
| | Human | MAB8692 (ICC/IF) | | | |
| ◆ p38β | Human | MAB3274 (ICC/IF) | | | ✓ |
| ◆ p38γ | Human/Mouse/Rat | AF1347 (IHC, SW, WB) | | | ✓ |
| | Human/Mouse/Rat | MAB1347 (SW, WB) | | | |
| | Human/Mouse/Rat | AF1644 (WB) | | | |
| ◆ p38δ | Human | AF1519 (IHC, WB) | | | ✓ |
| | Human | MAB1519 (WB) | | | |
| ◆ c-Rel | Human | MAB4606 (ICC/IF, WB) | | ARY029 | |
| | Human/Mouse | AF2699 (ChIP, ICC/IF, SW, WB) | | | |
| | Mouse | MAB2699 (WB) | | | |

* Indicates a recombinant monoclonal antibody
Application key: B/N Blocking/Neutralization ChIP Chromatin Immunoprecipitation FA Functional Assay FC Flow Cytometry ICC/IF Immunocytochemistry/Immunofluorescence IHC Immunohistochemistry IP Immunoprecipitation SW Simple Western WB Western blot
Fluorochrome key for FAB/IC catalog numbers ending in: A Allophycocyanin C PerCP F Fluorescein G AlexaFluor® 488 N AlexaFluor® 700 P Phycoerythrin
◆ Indicates a R&D Systems brand antibody ◆ Indicates a Novus Biologicals brand antibody

| Molecule | R&D Systems™ or Novus Biologicals™ Antibodies | | R&D Systems™ ELISA/Activity Assay Kits Catalog # | R&D Systems™ Proteome Profiler™ Antibody Array Kits Catalog # | Tocris™ Small Molecule Activators/Inhibitors |
|--------------------------------|---|---------------------------|--|---|--|
| | Species | Catalog # (Applications) | | | |
| ◆ RelA/NFκB p65 | Human | MAB50781 (FC) | | ARY029 | |
| | Human/Mouse | MAB5078 (FC, ICC/IF, WB) | | | |
| | Human/Mouse | IC5078A, G, P (FC) | | | |
| | Human/Mouse | AF5078 (ChIP, SW, WB) | | | |
| ◆ Phospho-RelA/NFκB p65 (S529) | Human | MAB7624 (WB) | | ARY029 | |
| ◆ Phospho-RelA/NFκB p65 (S536) | Human | MAB7226 (ICC/IF, WB) | | | |
| | Human | MAB72261 (ICC/IF, SW, WB) | | | |
| ◆ RelB | Human | MAB2698 (ICC/IF, IHC, WB) | | | |
| ◆ TAB1 | Human/Mouse | AF3578 (ICC/IF, WB) | | | |
| ◆ TAK1 | Human | MAB5307 (WB) | | | ✓ |
| ◆ TRAF-6 | Human | AF3284 (WB) | | ARY027 | |

* Indicates a recombinant monoclonal antibody
Application key: B/N Blocking/Neutralization ChIP Chromatin Immunoprecipitation FA Functional Assay FC Flow Cytometry ICC/IF Immunocytochemistry/Immunofluorescence IHC Immunohistochemistry IP Immunoprecipitation SW Simple Western WB Western blot
Fluorochrome key for FAB/IC catalog numbers ending in: A Allophycocyanin C PerCP F Fluorescein G AlexaFluor® 488 N AlexaFluor® 700 P Phycoerythrin
◆ Indicates a R&D Systems brand antibody ◆ Indicates a Novus Biologicals brand antibody



Learn More

Scan the QR Code or visit:

[bio-techne.com/il-1family](https://www.bio-techne.com/il-1family)

Tocris™ IL-1 Signaling-related Small Molecule Activators & Inhibitors

Tocris is the leading supplier of high performance reagents for life science research. The Tocris range of small molecules and peptides includes novel and exclusive cytokine receptor antagonists along with activators and inhibitors of specific kinases and transcription factors that regulate cytokine signaling pathways, including IL-1 family cytokine signaling.

| Category | Product Name | Description | Cat.# |
|--|-------------------|---|-------|
| IL-1 Family Cytokines & Receptors | | | |
| IL-1 Inhibitors | AF 12198 | Potent, selective human type I IL-1 receptor antagonist | 1793 |
| | CRID3 sodium salt | Potent NLRP3 inflammasome inhibitor; inhibits IL-1 β production | 5479 |
| IL-1 Family Cytokine Signaling | | | |
| AP-1 Inhibitors | c-Jun Peptide | Peptide inhibitor of JNK/c-Jun interaction | 1989 |
| | SR 11302 | AP-1 inhibitor | 2476 |
| ERK Inhibitors | FR 180204 | Selective ERK inhibitor | 3706 |
| | TCS ERK 11e | Potent and selective ERK2 inhibitor | 4465 |
| IKK Inhibitors | ACHP | Selective IKK α and IKK β inhibitor | 4547 |
| | BMS 345541 | Selective allosteric inhibitor of IKK; anti-inflammatory | 4806 |
| | IKK 16 | Selective inhibitor of IKK | 2539 |
| | IMD 0354 | Inhibitor of IKK β | 2611 |
| | ML 120B | IKK2-selective inhibitor | 4899 |
| | PF 184 | Potent and selective IKK β inhibitor | 4238 |
| | TPCA-1 | Potent, selective inhibitor of IKK β | 2559 |

| Category | Product Name | Description | Cat.# |
|-----------------------------------|---------------------|---|-------|
| IRAK Inhibitors | AS 2444697 | Potent and selective IRAK4 inhibitor | 5430 |
| | IRAK1/4 Inhibitor I | IRAK1 and IRAK4 inhibitor | 5665 |
| JNK Activators/ Inhibitors | Anisomycin | JNK, SAPK, and p38 activator | 1290 |
| | CEP 1347 | Inhibitor of JNK signaling | 4924 |
| | SP 600125 | Selective JNK inhibitor | 1496 |
| | TCS JNK 5 α | Selective inhibitor of JNK2 and JNK3 | 2827 |
| | TCS JNK 6 α | Selective JNK inhibitor | 3222 |
| p38 MAPK Inhibitors | AMG 548 | Potent and selective p38 α inhibitor | 3920 |
| | RWJ 67657 | Potent and selective p38 α and p38 β inhibitor | 2999 |
| | SB 202190 | Potent and selective p38 MAPK inhibitor | 1264 |
| | SB 203580 | Selective inhibitor of p38 MAPK; water soluble | 1402 |
| | SB 239063 | Potent and selective p38 MAPK inhibitor | 1962 |
| | SCIO 469 | Selective p38 MAPK inhibitor | 3528 |
| TAK1 Inhibitors | (5Z)-7-Oxozeaenol | Potent and selective TAK1 MAP3K inhibitor | 3604 |

| Category | Product Name | Description | Cat.# |
|--|-----------------------------------|---|-------|
| IκB/NFκB Activators/ Inhibitors | Betulinic acid | Activates NF κ B; anti-tumor and anti-HIV agent | 3906 |
| | Arctigenin | Inhibitor of I κ B α phosphorylation; also inhibits MEK1 | 1777 |
| | Bay 11-7821 | Indirect inhibitor of I κ B α phosphorylation | 1744 |
| | Caffeic acid phenethyl ester | Specific inhibitor of NF κ B activation | 2743 |
| | Cardamonin | Inhibitor of NF κ B activation; anti-inflammatory | 2509 |
| | Honokiol | Blocks NF κ B activation; also anti-inflammatory and antioxidant | 4590 |
| | IP7e | Blocks NF κ B pathway | 5699 |
| | Luteolin | Blocks NF κ B activation | 2874 |
| | MG 132 | Inhibits NF κ B activation; proteasome and calpain inhibitor | 1748 |
| | SP 100030 | NF κ B and AP-1 dual inhibitor | 5309 |
| | Sulfasalazine | Inhibitor of NF κ B activation | 4935 |
| Withaferin A | Inhibits NF κ B activation | 2816 | |



Learn More

Scan the QR Code or visit:

bio-techne.com/reagents/small-molecules-and-peptides

Contact Us

Global info@bio-techne.com, bio-techne.com/find-us/distributors

North America TEL 800 343 7475

Europe // Middle East // Africa TEL +44 (0)1235 529449

China info.cn@bio-techne.com, TEL 400.821.3475

For research use or manufacturing purposes only. Trademarks and registered trademarks are the property of their respective owners.
7486902496_1124

bio-techne[®]

Global Developer, Manufacturer, and Supplier of High-Quality Reagents,
Analytical Instruments, and Precision Diagnostics.

INCLUDES R&D Systems™ Novus Biologicals™ Tocris Bioscience™ ProteinSimple™ ACD™ ExosomeDx™ Asuragen™ Lunaphore™