

DESCRIPTION

Source	Porcine platelet-derived
Structure / Form	Disulfide-linked homodimer

SPECIFICATIONS

SDS-PAGE	12 kDa, reducing conditions
Activity	Measured by its ability to inhibit the IL-4-dependent proliferation of HT-2 mouse T cells. Tsang, M. <i>et al.</i> (1995) Cytokine 7:389. The ED ₅₀ for this effect is 0.03-0.2 ng/mL.
Endotoxin Level	<0.10 EU per 1 µg of the protein by the LAL method.
Purity	>97%, by SDS-PAGE under reducing conditions and visualized by silver stain.
Formulation	Lyophilized from a 0.2 µm filtered solution in Acetonitrile and TFA. See Certificate of Analysis for details.

PREPARATION AND STORAGE

Reconstitution	Reconstitute 1 µg vials at 10 µg/mL in sterile 4 mM HCl containing at least 0.1% human or bovine serum albumin. Reconstitute 10 µg or larger vials at 100 µg/mL in sterile 4 mM HCl.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 3 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

TGF-β1 (transforming growth factor beta 1) is one of three closely related mammalian members of the large TGF-β superfamily that share a characteristic cystine knot structure (1 - 7). TGF-β1, -2 and -3 are highly pleiotropic cytokines that are proposed to act as cellular switches that regulate processes such as immune function, proliferation and epithelial-mesenchymal transition (1 - 4). Each TGF-β isoform has some non redundant functions; for TGF-β1, mice with targeted deletion show defects in hematopoiesis and endothelial differentiation, and die of overwhelming inflammation (2). Porcine TGF-β1 cDNA encodes a 390 amino acid (aa) precursor that contains a 29 aa signal peptide and a 361 aa proprotein (8). A furin-like convertase processes the proprotein to generate an N-terminal 220 aa latency-associated peptide (LAP) and a C-terminal 112 aa mature TGF-β1 (8, 9). Disulfide-linked homodimers of LAP and TGF-β1 remain non-covalently associated after secretion, forming the small latent TGF-β1 complex (8 - 10). Covalent linkage of LAP to one of three latent TGF-β binding proteins (LTBPs) creates a large latent complex that may interact with the extracellular matrix (9, 10). TGF-β is activated from latency by pathways that include actions of the protease plasmin, matrix metalloproteases, thrombospondin 1 and a subset of integrins (10). Mature porcine TGF-β1 shows 100% aa identity with human, dog and cow TGF-β1 and 99% aa identity with mouse, rat and horse TGF-β1. It demonstrates cross species activity (1). TGF-β1 signaling begins with high-affinity binding to a type II ser/thr kinase receptor termed TGF-β RII. This receptor then phosphorylates and activates a second ser/thr kinase receptor, TGF-β RI (also called activin receptor-like kinase (ALK) -5), or alternatively, ALK-1. This complex phosphorylates and activates Smad proteins that regulate transcription (3, 11, 12). Contributions of the accessory receptors betaglycan (also known as TGF-β RIII) and endoglin, or use of Smad-independent signaling pathways, allow for disparate actions observed in response to TGF-β in different contexts (11).

References:

1. Sporn, M.B. (2006) Cytokine Growth Factor Rev. 17:3.
2. Dunker, N. and K. Kriegelstein (2000) Eur. J. Biochem. 267:6982.
3. Wahl, S.M. (2006) Immunol. Rev. 213:213.
4. Chang, H. *et al.* (2002) Endocr. Rev. 23:787.
5. Lin, J.S. *et al.* (2006) Reproduction 132:179.
6. Hinck, A.P. *et al.* (1996) Biochemistry 35:8517.
7. Mittl, P.R.E. *et al.* (1996) Protein Sci. 5:1261.
8. Kondaiah, P. *et al.* (1988) J. Biol. Chem. 263:18313.
9. Miyazono, K. *et al.* (1988) J. Biol. Chem. 263:6407.
10. Oklu, R. and R. Hesketh (2000) Biochem. J. 352:601.
11. De Caestecker, M. *et al.* (2004) Cytokine Growth Factor Rev. 15:1.
12. Zuniga, J.E. *et al.* (2005) J. Mol. Biol. 354:1052.v. 15:1.