biotechne® RDSYSTEMS

Recombinant Human Melanotransferrin/CD228 His-tag

Catalog Number: 10696-MT

DESCRIPTION	
Source	Human embryonic kidney cell, HEK293-derived human Melanotransferrin/CD228 protein Gly20-Gly711, with a C-terminal 6-His tag Accession # NP_005920.2
N-terminal Sequence Analysis	Gly20
Predicted Molecular Mass	76 kDa

SPECIFICATIONS	
SDS-PAGE	84-92 kDa, under reducing conditions
Activity	Measured by its binding ability in a functional ELISA. Recombinant Human Melanotransferrin/CD228 His-tag (Catalog # 10696-MT) binds Human Melanotransferrin/CD228 Antibody (Catalog # MAB81751) with an ED ₅₀ of < 400 ng/mL.
Endotoxin Level	<0.10 EU per 1 μ g of the protein by the LAL method.
Purity	>95%, by SDS-PAGE visualized with Silver Staining and quantitative densitometry by Coomassie® Blue Staining.
Formulation	Lyophilized from a 0.2 μm filtered solution in PBS. See Certificate of Analysis for details.

PREPARATION AND	STORAGE
Reconstitution	Reconstitute at 500 μg/mL in PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.
	 12 months from date of receipt, -20 to -70 °C as supplied.
	1 month 2 to 8 °C under sterile conditions after reconstitution

• 3 months, -20 to -70 °C under sterile conditions after reconstitution.

DATA			
SDS-PAGE			
	kDa R 190 - 92 - 66 - 55 - 43 - 29 - 21 - 18 - 12 - 6 -	NR	Recombinant Human Melanotransferrin/CD228 His- tag Protein SDS PAGE 2 µg/lane of Recombinant Human Melanotransferrin/CD228 His-tag Protein (Catalog # 10696-MT) was resolved with SDS-PAGE under reducing (R) and non- reducing (NR) conditions and visualized by Coomassie® Blue staining, showing bands at 84-92 kDa and 75-85 kDa, respectively.

BACKGROUND

Melanotransferrin (MTF), also known as CD228, P97, MELTF, or MFI2, is a member of the transferrin superfamily that binds to a single ferric iron with high affinity (1). It was originally identified as a membrane-bound protein with a coding region of 2217 bases encoding 738 amino acids (aa) including five Fe-binding residues, the consensus thermolysin metalloprotease sites, and a glycosyl-phosphatidylinositol (GPI) anchor (2). A secreted form of MTF (sMTF) has also been reported (2, 3). Human MTF shares 86% as sequence identity with mouse and rat MTF. MTF expression is low in normal tissues but high in tumor and embryonic tissues. In melanoma MTF is associated with tumor metastasis and angiogenesis and upregulated in lung cancer tissue and cell lines (3). In the brain MTF is expressed in capillary endothelium but also in the reactive microglia associated with senile plaques in Alzheimer's disease (AD) (4). sMTF has been reported to be increased in patients with AD or arthritis and may modulate angiogenesis, cell migration, and plasminogen activation (5).

References:

- 1. Baker, E.N. *et al*. (1992) FEBS Lett. **298**:215.
- 2. Sekyere, E. & D.R. Richardson (2000) FEBS Lett. 483:11.
- 3. Lei, Y. et al. (2020) Cell Death Dis. 11:933.
- 4. Jefferies, W.A. et al. (1996) Brain Res. 712:122.
- 5. Paluncic, J. et al. (2016) Biochim. Biophy. Acta. 1863:770.

Rev. 4/28/2025 Page 1 of 1

Bio-Techne[®] Global | bio-techne.com info@bio-techne.com techsupport@bio-techne.com TEL: 1.612.379.2956 USA | TEL: 800.343.7475 Canada | TEL: 855.668.8722 Europe | Middle East | Africa TEL: +44.0.1235.529449 China | info.cn@bio-techne.com TEL: 400.821.3475