

Recombinant HCoV-OC43 Spike His-tag

Catalog Number: 10989-CV

DESCRIPTION	
Source	Human embryonic kidney cell, HEK293-derived hcov-oc43 Spike protein Val15-Pro1297 (Arg754Ser, Arg757Ser), with a C-terminal 6-His tag Accession # YP_009555241.1
N-terminal Sequence Analysis	Val15
Predicted Molecular Mass	144 kDa

SPECIFICATIONS	
SDS-PAGE	>150 kDa, under reducing conditions.
Activity	Measured by its ability to agglutinate mouse red blood cells. The ED_{50} for this effect is 0.10-1.00 µg/mL.
Endotoxin Level	<0.10 EU per 1 µg of the protein by the LAL method.
Purity	>95%, by SDS-PAGE visualized with Silver Staining and quantitative densitometry by Coomassie® Blue Staining.
Formulation	Supplied as a 0.2 µm filtered solution in PBS. See Certificate of Analysis for details.

PREPARATION AND STORAGE		
Shipping	The product is shipped with dry ice or equivalent. Upon receipt, store it immediately at the temperature recommended below.	
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.	
	 6 months from date of receipt, -20 to -70 °C as supplied. 	
	1 month, 2 to 8 °C under sterile conditions after opening.	
	 3 months -20 to -70 °C under sterile conditions after opening 	

DATA SDS-PAGE kDa NR Recombinant HCoV-OC43 Spike His-tag Protein SDS-PAGE. 2 µg/lane of Recombinant 190 HCoV-OC43 Spike His-tag Protein (Catalog # 10989-CV) was resolved with SDS-PAGE 92 under reducing (R) and non-66 reducing (NR) conditions and 55 -43 visualized by Coomassie® Blue staining, showing bands at >150 kDa. 29 -21 -12

BACKGROUND

HCoV-OC43, a virus first isolated in 1960's that accounts for ~ 20% of the common cold, belongs to a family of viruses known as coronaviruses that are commonly comprised of a large plus-strand RNA genome and four structural proteins: Spike protein (S), Envelope protein (E), Membrane protein (M), and Nucleocapsid protein (N) (1, 2). Other well-known human coronaviruses include three viruses that cause relatively mild respiratory disease: HCoV-229E, HCoV-HKU1 and HCov-NL63, plus three viruses that cause the Severe Acute Respiratory Syndrome (SARS-CoV), the Middle East Respirator Syndrome (MERS-CoV), and the global pandemic Covid-19 (SARS-CoV2). While the S, E and M proteins build up the viral envelop, the N protein is involved transcription, replication, and packaging of the viral RNA genome into a helical ribonucleocapsid (RNP) (3, 4). The CoV-OC43 N protein is a ~50 kDa protein composed of two independent structural domains connected by a linker region. Both the N-terminal and the linker regions contain RNA binding domains, while the C-terminal region is responsible for the oligomerization of the N protein (5). The CoV-OC43 N protein same definition and displays high immunogenic activity. Cross activity of antibodies among different strains should be rigorously tested when designing serological diagnostic kits (6, 7).

References:

- 1. St-Jean, J.R. *et al.* (2004) J. Virol. **78**:8824.
- 2. Vabret, A. et al. (2003) Fr Clin Infect Dis. 36:985.
- 3. Chang, C.K. et al. (2006) J. Biomed. Sci. 13:59.
- 4. Hurst, K.R. et al. (2009) J. Virol. 83:7221.
- 5. Huang, C.Y. et al. (2009) Protein Sci. 18:2209.
- 6. Chan, K.H. et al. (2005) Clin Diagn Lab Immunol. 12:1317.
- 7. Mourez, T. et al. (2007) J. Virol Methods. 139:175.

Rev. 8/25/2022 Page 1 of 1



Global bio-techne.com info@bio-techne.com techsupport@bio-techne.com TEL +1 612 379 2956 USA TEL 800 343 7475 **Canada** TEL 855 668 8722 **China** TEL +86 (21) 52380373 **Europe | Middle East | Africa** TEL +44 (0)1235 529449