

DESCRIPTION

Source	Human embryonic kidney cell, HEK293-derived sars-cov-2 Spike protein		
	SARS-CoV-2 R.1 Spike (Val16-Lys1211) (Trp152Leu, Glu484Lys, Asp614Gly, Gly769Val) (Arg682Ser, Arg685Ser, Lys986Pro, Val987Pro) Accession # YP_009724390.1	GCN4-IZ	6-His tag
	N-terminus		C-terminus
N-terminal Sequence Analysis	Val16		
Predicted Molecular Mass	138 kDa		

SPECIFICATIONS


SDS-PAGE	154 - 174 kDa, under reducing conditions.
Activity	Measured by its binding ability in a functional ELISA with Recombinant Human ACE-2 His-tag (Catalog # 933-ZN).
Endotoxin Level	<0.10 EU per 1 µg of the protein by the LAL method.
Purity	>95%, by SDS-PAGE visualized with Silver Staining and quantitative densitometry by Coomassie® Blue Staining.
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details.

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 500 µg/mL in PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> • 12 months from date of receipt, -20 to -70 °C as supplied. • 1 month, 2 to 8 °C under sterile conditions after reconstitution. • 3 months, -20 to -70 °C under sterile conditions after reconstitution.


DATA

Binding Activity



Recombinant SARS-CoV-2 R.1 S (GCN4-IZ) His-tag Protein Binding Activity.
Recombinant SARS-CoV-2 R.1 S (GCN4-IZ) His-tag (Catalog # 11026-CV) binds Recombinant Human ACE-2 His-tag (Catalog # 933-ZN) in a functional ELISA.

SDS-PAGE



Recombinant SARS-CoV-2 R.1 S (GCN4-IZ) His-tag Protein SDS-PAGE. 2 µg/lane of Recombinant SARS-CoV-2 R.1 S (GCN4-IZ) His-tag Protein (Catalog # 11026-CV) was resolved with SDS-PAGE under reducing (R) and non-reducing (NR) conditions and visualized by Coomassie® Blue staining, showing bands at 154-174 kDa.

BACKGROUND

SARS-CoV-2, which causes the global pandemic coronavirus disease 2019 (Covid-19), belongs to a family of viruses known as coronaviruses that also include MERS and SARS-CoV-1. Coronaviruses are commonly comprised of four structural proteins: Spike protein (S), Envelope protein (E), Membrane protein (M) and Nucleocapsid protein (N) (1). The SARS-CoV-2 S protein is a glycoprotein that mediates membrane fusion and viral entry. The S protein is homotrimeric, with each ~180 kDa monomer consisting of two subunits, S1 and S2 (2). In SARS-CoV-2, as with most coronaviruses, proteolytic cleavage of the S protein into S1 and S2 subunits is required for activation. The S1 subunit is focused on attachment of the protein to the host receptor while the S2 subunit is involved with cell fusion (3-5). A metalloproteinase, angiotensin-converting enzyme 2 (ACE-2), has been identified as a functional receptor for SARS-CoV-2 through interaction with a receptor binding domain (RBD) located at the C-terminus of S1 subunit (6, 7). The S protein of SARS-CoV-2 shares 75% and 29% amino acid (aa) sequence identity with the S protein of SARS-CoV-1 and MERS, respectively. A SARS-CoV-2 variant (R.1) carrying the aa substitution Trp152Leu, Glu484Lys, Asp614Gly, and Gly769Val in the spike protein was first identified in January 2021 in Japan (8). Whether these mutations in the spike would cause more severe symptom or decrease the efficacy of vaccine-induced immunity is still under investigation.

References:

1. Wu, F. *et al.* (2020) *Nature* **579**:265.
2. Tortorici, M.A. and D. Veasley (2019) *Adv. Virus Res.* **105**:93.
3. Bosch, B.J. *et al.* (2003) *J. Virol.* **77**:8801.
4. Belouzard, S. *et al.* (2009) *Proc. Natl. Acad. Sci.* **106**:5871.
5. Millet, J.K. and G.R. Whittaker (2015) *Virus Res.* **202**:120.
6. Li, W. *et al.* (2003) *Nature* **426**:450.
7. Wong, S.K. *et al.* (2004) *J. Biol. Chem.* **279**:3197.
8. Hirotsu, Y. and M. Omata (2021) *PLoS Pathog.* **17**:e1009619.