

DESCRIPTION

Source	Mouse myeloma cell line, NS0-derived			
	Mouse Netrin-1 (Val22-Ala603) Accession # AAC52971 N-terminus	IEGR	GGGSGGGSGGGS	10-His tag C-terminus

N-terminal Sequence Val22

Analysis

Predicted Molecular Mass 68.2 kDa


SPECIFICATIONS

SDS-PAGE	83 kDa, reducing conditions
Activity	Measured by its binding ability in a functional ELISA. Immobilized recombinant rat UNC5H2 Fc Chimera at 5 µg/mL (100 µL/well) can bind Recombinant Mouse Netrin-1 with a linear range of 3-200 ng/mL.
Endotoxin Level	<0.10 EU per 1 µg of the protein by the LAL method.
Purity	>90%, by SDS-PAGE visualized with Silver Staining and quantitative densitometry by Coomassie® Blue Staining.
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS and EDTA with Trehalose. See Certificate of Analysis for details.

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 100 µg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	<p>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</p> <ul style="list-style-type: none"> • 12 months from date of receipt, -20 to -70 °C as supplied. • 1 month, 2 to 8 °C under sterile conditions after reconstitution. • 3 months, -20 to -70 °C under sterile conditions after reconstitution.

DATA



BACKGROUND

Mouse Netrin-1 is a member of the laminin-related family of axon-guidance molecules, collectively referred to as Netrins (*netr* is Sanskrit for "one who guides"). The molecule's cDNA encodes a 603 amino acid (aa) protein precursor that has structural similarity to the N-terminal γ -chain of laminin. It contains a globular domain, three EGF repeats, and a C-terminal heparin-binding domain. Mouse Netrin-1 shares 52% aa identity with mouse Netrin-3, and 98% and 87% aa identity with human and chicken Netrin-1, respectively. Cells reported to express Netrin-1 in the embryo include cells of the floor plate, ventricular zone of the spinal cord, the brain, the ganglionic eminence, and parts of the diencephalon. Netrins were first identified for promoting the outgrowth of commissural axons and are also involved in helping migrating cells and axonal growth cones navigate to their targets. Netrins can provide both attractive and repulsive cues to neurons, depending on the receptors present and cellular context. In the adult, Netrin-1 is likely involved in axon regeneration in peripheral nerves. Netrin-1 has also been shown to be expressed outside of the nervous system and to be involved in development of such tissues as the pancreas, lung, bowel, bone and mammary gland. In non-neural organogenesis, Netrin-1 provides an adhesive rather than guidance function. The DCC (deleted in colorectal carcinoma), Neogenin, the UNC5 family of receptors, and the adenosine A2b receptors are proposed to be functional receptors for Netrin-1 (1-7).

References:

1. Puschel, A. (1999) *Mech. Dev.* **83**:65.
2. Hedgecock, E. and C. Norris (1997) *Trends Genet.* **13**:251.
3. Kappler, J. *et al.* (2000) *Biochem. Biophys. Res. Commun.* **271**:287.
4. Madison, R. *et al.* (2000) *Exp. Neurology* **161**:563.
5. Srinivasan, K. *et al.* (2003) *Dev. Cell* **4**:371.
6. Livesey, F.J. (1999) *Cell Mol. Life Sci.* **56**:62.
7. Corset, V. *et al.* (2000) *Nature* **407**:747.

PRODUCT SPECIFIC NOTICES

This product or the use of this product is covered by U.S. Patents owned by The Regents of the University of California. This product is for research use only and is not to be used for commercial purposes. Use of this product to produce products for sale or for diagnostic, therapeutic or drug discovery purposes is prohibited. In order to obtain a license to use this product for such purposes, contact The Regents of the University of California.

U.S. Patent # 5,565,331, 6,096,866, 6,017,714, 6,309,638, 6,670,451, and other U.S. and international patents pending.