

DESCRIPTION

Source Human embryonic kidney cell, HEK293-derived human KGF/FGF-7 protein
Cys32-Thr194
Accession # P21781.1

N-terminal Sequence Analysis Cys32

Predicted Molecular Mass 19 kDa

SPECIFICATIONS

SDS-PAGE 22-28 kDa, under reducing conditions.

Activity Measured in a serum-free cell proliferation assay using MCF-7 human breast cancer cells. Karey, K.P. *et al.* (1988) Cancer Research 48:4083.
The ED₅₀ for this effect is 25.0-250 ng/mL.

Endotoxin Level <0.10 EU per 1 µg of the protein by the LAL method.

Purity >95%, by SDS-PAGE visualized with Silver Staining and quantitative densitometry by Coomassie® Blue Staining.

Formulation Lyophilized from a 0.2 µm filtered solution in MOPS, Na₂SO₄ and EDTA with Trehalose. See Certificate of Analysis for details.

PREPARATION AND STORAGE

Reconstitution Reconstitute 10 µg size at 100 µg/mL and all the other sizes at 250 µg/mL in PBS.


Shipping The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

- 12 months from date of receipt, -20 to -70 °C as supplied.
- 1 month, 2 to 8 °C under sterile conditions after reconstitution.
- 3 months, -20 to -70 °C under sterile conditions after reconstitution.


DATA

Bioactivity



Recombinant Human KGF/FGF-7 Protein Bioactivity. Recombinant Human KGF/FGF-7 Protein (Catalog # 11340-KG) stimulates cell proliferation of MCF-7 human breast cancer cells. The ED₅₀ for this effect is 25.0-250 ng/mL.

SDS-PAGE



Recombinant Human KGF/FGF-7 Protein SDS-PAGE. 2 µg/lane of Recombinant Human KGF/FGF-7 Protein (Catalog # 11340-KG) was resolved with SDS-PAGE under reducing (R) and non-reducing (NR) conditions and visualized by Coomassie® Blue staining, showing bands at 22-28 kDa.

BACKGROUND

KGF (keratinocyte growth factor), also known as FGF-7 (fibroblast growth factor-7), is one of 22 known members of the mouse FGF family of secreted proteins that plays a key role in development, morphogenesis, angiogenesis, wound healing, and tumorigenesis (1-4,13). KGF expression is restricted to cells of mesenchymal origin. When secreted, it acts as a paracrine growth factor for nearby epithelial cells (1). KGF speeds wound healing by being dramatically upregulated in response to damage to skin or internal structures that results in high local concentrations of inflammatory mediators such as IL-1 and TNF-alpha. (2, 5). KGF promotes cell migration and invasion, and mediates melanocyte transfer to keratinocytes upon UVB radiation (6, 7). It has been used ectopically to avoid chemotherapy-induced oral mucositis in patients with hematological malignancies (1). Deletion of KGF affects kidney development, producing abnormally small ureteric buds and fewer nephrons (8). It also impedes hair follicle differentiation (9). The 194 amino acid (aa) KGF precursor contains a 31 aa signal sequence and, like all other FGFs, an ~120 aa beta -trefoil scaffold that includes receptor- and heparin-binding sites. KGF signals only through the IIIb splice form of the tyrosine kinase receptor, FGF R2 (FGF R2-IIIb/KGF R) (10). Receptor dimerization requires an octameric or larger heparin or heparin sulfate proteoglycan (11). FGF-10, also called KGF2, shares 51% aa identity and similar function to KGF, but shows more limited expression than KGF and uses an additional receptor, FGF R2-IIIc (12). Following receptor engagement, KGF is typically degraded, while FGF-10 is recycled (12). Mature human KGF, which is active across species, shares 98% aa sequence identity with bovine, equine, ovine and canine, 96% with mouse and porcine, and 92% with rat KGF, respectively.

References:

1. Finch, P.W. and J.S. Rubin (2006) *J. Natl. Cancer Inst.* **98**:812.
2. Werner, S. *et al.* (2007) *J. Invest. Dermatol.* **127**:998.
3. Werner, S. (1998) *Cytokine Growth Factor Rev.* **9**:153.
4. Mason, I.J. *et al.* (1994) *Mech. Dev.* **45**:15.
5. Geer, D.J. *et al.* (2005) *Am. J. Pathol.* **167**:1575.
6. Niu, J. *et al.* (2007) *J. Biol. Chem.* **282**:6001.
7. Cardinali, G. *et al.* (2005) *J. Invest. Dermatol.* **125**:1190.
8. Qiao, J. *et al.* (1999) *Development* **126**:547.
9. Guo, L. *et al.* (1996) *Genes Dev.* **10**:165.
10. de Georgi, V. *et al.* (2007) *Dermatol. Clin.* **25**:477.
11. Hsu, Y-R. *et al.* (1999) *Biochemistry* **38**:2523.
12. Belleudi, F. *et al.* (2007) *Traffic* **8**:1854.
13. Vadija R. *et al.* (2016) *J Chem Biol.* **9**:69.