

#### DESCRIPTION

Source	Chinese Hamster Ovary cell line, CHO-derived human TSPAN13 protein			
	MD	Human IgG <sub>1</sub> (Pro100-Lys330)	IEGR	Human TSPAN13-LEL (Cys94-Arg167) Accession # O95857.1
	N-terminus		C-terminus	
N-terminal Sequence Analysis	Met			
Structure / Form	Disulfide-linked homodimer			
Predicted Molecular Mass	35 kDa			

#### SPECIFICATIONS


<b>SDS-PAGE</b>	40-58 kDa, under reducing conditions
<b>Activity</b>	Bioassay data are not available.
<b>Endotoxin Level</b>	<0.10 EU per 1 µg of the protein by the LAL method.
<b>Purity</b>	>95%, by SDS-PAGE visualized with Silver Staining and quantitative densitometry by Coomassie® Blue Staining.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details.

#### PREPARATION AND STORAGE

<b>Reconstitution</b>	Reconstitute at 500 µg/mL in PBS.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
<b>Stability &amp; Storage</b>	<p><b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b></p> <ul style="list-style-type: none"> <li>12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>3 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

#### DATA

##### SDS-PAGE



**Recombinant Human TSPAN13-LEL Fc Chimera Protein SDS-PAGE.** 2 µg/lane of Recombinant Human TSPAN13-LEL Fc Chimera Protein (Catalog # 11551-TS) was resolved with SDS-PAGE under reducing (R) and non-reducing (NR) conditions and visualized by Coomassie® Blue staining, showing bands at 40-58 kDa and 80-120 kDa, respectively.

#### BACKGROUND

Tetraspanin 13, or TSPAN13, belongs to a superfamily of proteins that is characterized by four transmembrane domains, three intracellular domains and two extracellular loops: a small extracellular loop (SEL) and a large extracellular loop (LEL). The extracellular loops form molecular webs that bring together cell surface proteins, facilitating the formation of stable and functional signalling complexes. Tetraspanins form microdomains on the plasma membrane that mediate diverse biological processes including adhesion, cell fusion, immune response, and tumor development (1-4). Human TSPAN13 consists of 204 amino acids, with the LEL region spanning residues 94-167. Within the LEL, human TSPAN13 shares 89% aa 88% aa sequence identity with mouse and rat TSPAN13-LEL, respectively.

#### References:

1. Charrin, S. *et al.* (2014) J. Cell Sci. **127**:3641.
2. Yang, J. *et al.* (2024) Cell. **13**:193.
3. Hemler, M.E. (2005) Nat. Rev. Mol. Cell Biol. **6**:801.
4. Kim, T-K. *et al.* (2015) Biochem. Biophys. Res. Commun. **468**:774.