

DESCRIPTION

Source *E. coli*-derived
Pro29-Met212
Accession # Q75MH2

N-terminal Sequence Analysis Pro29

Predicted Molecular Mass 20.3 kDa

SPECIFICATIONS

Activity Measured in a cell proliferation assay using T1165.85.2.1 mouse plasmacytoma cells. Nordan, R.P. *et al.* (1987) J. Immunol. **139**:813. The ED₅₀ for this effect is 0.2-0.8 ng/mL. The specific activity of Recombinant Human IL-6 is approximately 1.1 x 10⁵ IU/μg, which is calibrated against human IL-6 WHO International Standard (NIBSC code: 89/548).

Endotoxin Level <0.10 EU per 1 μg of the protein by the LAL method.

Purity >97%, by SDS-PAGE under reducing conditions and visualized by silver stain.

Formulation Lyophilized from a 0.2 μm filtered solution in PBS and NaCl with BSA as a carrier protein. See Certificate of Analysis for details.

PREPARATION AND STORAGE



Reconstitution Reconstitute at 100-200 μg/mL in sterile PBS containing at least 0.1% human or bovine serum albumin.

Shipping The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

- 12 months from date of receipt, -20 to -70 °C as supplied.
- 1 month, 2 to 8 °C under sterile conditions after reconstitution.
- 3 months, -20 to -70 °C under sterile conditions after reconstitution.

DATA



BACKGROUND

Interleukin-6 (IL-6) is a pleiotropic, α -helical, 22-28 kDa phosphorylated and variably glycosylated cytokine that plays important roles in the acute phase reaction, inflammation, hematopoiesis, bone metabolism, and cancer progression (1-5). Mature human IL-6 is 183 amino acids (aa) in length and shares 39% aa sequence identity with mouse and rat IL-6 (6). Alternative splicing generates several isoforms with internal deletions, some of which exhibit antagonistic properties (7-10). IL-6 induces signaling through a cell surface heterodimeric receptor complex composed of a ligand binding subunit (IL-6 R α) and a signal transducing subunit (gp130). IL-6 binds to IL-6 R α , triggering IL-6 R α association with gp130 and gp130 dimerization (11). gp130 is also a component of the receptors for CLC, CNTF, CT-1, IL-11, IL-27, LIF, and OSM (12). Soluble forms of IL-6 R α are generated by both alternative splicing and proteolytic cleavage (5). In a mechanism known as trans-signaling, complexes of soluble IL-6 and IL-6 R α elicit responses from gp130-expressing cells that lack cell surface IL-6 R α (5). Trans-signaling enables a wider range of cell types to respond to IL-6, as the expression of gp130 is ubiquitous, while that of IL-6 R α is predominantly restricted to hepatocytes, monocytes, and resting lymphocytes (2, 5). Soluble splice forms of gp130 block trans-signaling from IL-6/IL-6 R α but not from other cytokines that use gp130 as a co-receptor (5, 13). IL-6, along with TNF- α and IL-1, drives the acute inflammatory response and the transition from acute inflammation to either acquired immunity or chronic inflammatory disease (1-5). When dysregulated, it contributes to chronic inflammation in obesity, insulin resistance, inflammatory bowel disease, arthritis, sepsis, and atherosclerosis (1, 2, 5). IL-6 can also function as an anti-inflammatory molecule, as in skeletal muscle where it is secreted in response to exercise (2). In addition, it enhances hematopoietic stem cell proliferation and the differentiation of Th17 cells, memory B cells, and plasma cells (1, 14).

References:

1. Mansell, A. and B.J. Jenkins (2013) Cytokine Growth Factor Rev. **24**:249.
2. Schuett, H. *et al.* (2009) Thromb. Haemost. **102**:215.
3. Erta, M. *et al.* (2012) Int. J. Biol. Sci. **8**:1254.
4. Garbers, C. *et al.* (2012) Cytokine Growth Factor Rev. **23**:85.
5. Mihara, M. *et al.* (2012) Clin. Sci. (Lond.) **122**:143.
6. Hirano, T. *et al.* (1986) Nature **324**:73.
7. Kestler, D.P. *et al.* (1995) Blood **86**:4559.
8. Kestler, D.P. *et al.* (1999) Am. J. Hematol. **61**:169.
9. Bihl, M.P. *et al.* (2002) Am. J. Respir. Cell Mol. Biol. **27**:48.
10. Alberti, L. *et al.* (2005) Cancer Res. **65**:2.
11. Murakami, M. *et al.* (1993) Science **260**:1808.
12. Muller-Newen, G. (2003) Sci. STKE **2003**:PE40.
13. Mitsuyama, K. *et al.* (2006) Clin. Exp. Immunol. **143**:125.
14. Cerutti, A. *et al.* (1998) J. Immunol. **160**:2145.