

DESCRIPTION

Source Mouse myeloma cell line, NS0-derived mouse LYVE-1 protein
Ala24-Thr234, with a C-terminal 6-His tag
Accession # Q8BHC0

N-terminal Sequence Analysis Ala24

Predicted Molecular Mass 23.4 kDa

SPECIFICATIONS

SDS-PAGE 43-58 kDa, reducing conditions

Activity Measured by its ability to bind hyaluronan in a functional ELISA.

Endotoxin Level <0.10 EU per 1 µg of the protein by the LAL method.

Purity >95%, by SDS-PAGE visualized with Silver Staining and quantitative densitometry by Coomassie® Blue Staining.

Formulation Lyophilized from a 0.2 µm filtered solution in PBS. See Certificate of Analysis for details.

PREPARATION AND STORAGE

Reconstitution Reconstitute at 100 µg/mL in sterile PBS.

Shipping The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage **Use a manual defrost freezer and avoid repeated freeze-thaw cycles.**

- 12 months from date of receipt, -20 to -70 °C as supplied.
- 1 month, 2 to 8 °C under sterile conditions after reconstitution.
- 3 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Lymphatic vessel endothelial hyaluronan (HA) receptor-1 (LYVE-1) is a recently identified receptor of HA, a linear high molecular weight polymer composed of alternating units of D-glucuronic acid and N-acetyl-D-glucosamine. HA is found in the extracellular matrix of most animal tissues and in body fluids. It modulates cell behavior and functions during tissue remodeling, development, homeostasis, and disease (1). The turnover of HA (several grams/day in humans) occurs primarily in the lymphatics and liver, the two major clearance systems that catabolize approximately 85% and 15% of HA, respectively (1-3). LYVE-1 shares 41% homology with the other known HA receptor, CD44 (4). The homology between the two proteins increases to 61% within the HA binding domain. The HA binding domain, known as the link module, is a common structural motif found in other HA binding proteins such as link protein, aggrecan and versican (1, 5). Human and mouse LYVE-1 share 69% amino acid sequence identity.

LYVE-1 is primarily expressed on both the luminal and abluminal surfaces of lymphatic vessels (4, 5). In addition, LYVE-1 is also present in normal hepatic blood sinusoidal endothelial cells (6). LYVE-1 mediates the endocytosis of HA and may transport HA from tissue to lymph by transcytosis, delivering HA to lymphatic capillaries for removal and degradation in the regional lymph nodes (5, 7, 8). Because of its restricted expression patterns, LYVE-1, along with other lymphatic proteins such as VEGF R3, podoplanin and the homeobox protein prospero-related (Prox-1), constitute a set of markers useful for distinguishing between lymphatic and blood microvasculature (4, 5, 9-11).

References:

1. Knudson, C.B. and W. Knudson (1993) FASEB J. **7**:1233.
2. Evered, D. and J. Whelan (1989) Ciba Found. Symp. **143**:1.
3. Laurent, T.C. and J.R.F. Fraser (1992) FASEB J. **6**:2397.
4. Banerji, S. *et al.* (1999) J. Cell Biol. **144**:789.
5. Prevo, R. *et al.* (2001) J. Biol. Chem. **276**:19420.
6. Carreira, C.M. *et al.* (2001) Cancer Research **61**:8079.
7. Jackson, D.J. *et al.* (2001) Trends Immunol. **22**:317.
8. Zhou, B. *et al.* (2000) J. Biol. Chem. **275**:37733.
9. Achen, M. *et al.* (1998) Proc. Natl. Acad. Sci. USA **95**:548.
10. Breiteneder-Gellef, S. *et al.* (1999) Am. J. Pathol. **154**:385.
11. Wiggle, J.T. and G. Oliver (1999) Cell **98**:769.