

DESCRIPTION

Source	<i>Spodoptera frugiperda</i> , Sf 21 (baculovirus)-derived BDNF protein His129-Arg247 Accession # P23560
N-terminal Sequence Analysis	His129
Predicted Molecular Mass	13.5 kDa


SPECIFICATIONS

SDS-PAGE	13-14 kDa, reducing conditions
Activity	Measured in a cell proliferation assay using BaF mouse pro-B cells transfected with TrkB. The ED ₅₀ for this effect is 0.2-2 ng/mL. Measured by its binding ability in a functional ELISA. When Recombinant Human TrkB Fc Chimera (Catalog # 688-TK) is coated at 1 µg/mL, Recombinant Human BDNF binds with an apparent K _d <1 nM.
Endotoxin Level	<0.10 EU per 1 µg of the protein by the LAL method.
Purity	>97%, by SDS-PAGE visualized with Silver Staining and quantitative densitometry by Coomassie® Blue Staining.
Formulation	Lyophilized from a 0.2 µm filtered solution in Sodium Citrate and NaCl with BSA as a carrier protein. See Certificate of Analysis for details.

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 250 µg/mL in water.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	<ul style="list-style-type: none"> • 12 months from date of receipt, ≤ -20 °C as supplied. • 1 month, 2 to 8 °C under sterile conditions after reconstitution. • 3 months, ≤ -20 °C under sterile conditions after reconstitution.

DATA



BACKGROUND

Brain-derived neurotrophic factor (BDNF) is a member of the NGF family of neurotrophic factors (also named neurotrophins) that are required for the differentiation and survival of specific neuronal subpopulations in both the central as well as the peripheral nervous system. The neurotrophin family is comprised of at least four proteins including NGF, BDNF, NT-3, and NT-4/5. These secreted cytokines are synthesized as prepropeptides that are proteolytically processed to generate the mature proteins (1, 2). All neurotrophins have six conserved cysteine residues that are involved in the formation of three disulfide bonds and all share approximately 55% sequence identity at the amino acid level. Similarly to NGF, bioactive BDNF is predicted to be a non-covalently linked homodimer.

BDNF cDNA encodes a 247 amino acid residue precursor protein with a signal peptide and a proprotein that are cleaved to yield the 119 amino acid residue mature BDNF. The amino acid sequence of mature BDNF is identical in all mammals examined. High levels of expression of BDNF have been detected in the hippocampus, cerebellum, fetal eye and placenta. In addition, low levels of BDNF expression are also found in the pituitary gland, spinal cord, heart, lung and skeletal muscle. BDNF binds with high affinity and specifically activates the TrkB tyrosine kinase receptor (3).

References:

1. Eide, F.F. *et al.* (1993) *Exp. Neurol.* **121**:200.
2. Snider, W.D. (1994) *Cell* **77**:627.
3. Barbacid, M. (1994) *J. Neurobiol.* **25**:1386.