

DESCRIPTION

Source *E. coli*-derived human IFN-gamma protein
Gln24-Gln166 with an N-terminal Met
Accession # CAA31639

N-terminal Sequence Analysis Met

Predicted Molecular Mass 16.9 kDa

SPECIFICATIONS

SDS-PAGE 17 kDa, reducing conditions

Activity Measured in anti-viral assays using HeLa human cervical epithelial carcinoma cells infected with encephalomyocarditis (EMC) virus. Meager, A. (1987) in *Lymphokines and Interferons, a Practical Approach*. Clemens, M.J. *et al.* (eds): IRL Press. 129.
The ED₅₀ for this effect is 0.15-0.75 ng/mL.
The specific activity of Recombinant Human IFN- γ is approximately 2×10^4 IU/ μ g, which is calibrated against human IFN- γ Standard (NIBSC code: 87/586). Specific activity is for reference purposes only and is not routinely tested.

Endotoxin Level <0.01 EU per 1 μ g of the protein by the LAL method.

Purity >97%, by SDS-PAGE visualized with Silver Staining and quantitative densitometry by Coomassie® Blue Staining.

Formulation Lyophilized from a 0.2 μ m filtered solution in Sodium Succinate, Mannitol and Tween® 80. See Certificate of Analysis for details.

PREPARATION AND STORAGE

Reconstitution Reconstitute at 0.2 mg/mL in sterile, deionized water.


Shipping The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

- 12 months from date of receipt, -20 to -70 °C as supplied.
- 1 month, 2 to 8 °C under sterile conditions after reconstitution.
- 3 months, -20 to -70 °C under sterile conditions after reconstitution.


DATA

Bioactivity



Recombinant Human IFN-gamma Protein Bioactivity
Recombinant Human IFN- γ (Catalog # 285-IF/CF) demonstrates anti-viral activity in HeLa human cervical epithelial carcinoma cells infected with encephalomyocarditis (EMC) virus. The activity is over 2-fold greater than the top competitor's IFN- γ .

SDS-PAGE



Recombinant Human IFN-gamma Protein SDS-PAGE 1 μ g/lane of Recombinant Human IFN- γ was resolved with SDS-PAGE under reducing (R) conditions and visualized by silver staining, showing a single band at 17 kDa.

BACKGROUND

Interferon- γ (IFN- γ), also known as type II or immune interferon, exerts a wide range of immunoregulatory activities and is considered to be the prototype proinflammatory cytokine (1, 2). Mature human IFN- γ exists as a non-covalently linked homodimer of 20-25 kDa variably glycosylated subunits (3). It shares 90% amino acid (aa) sequence identity with rhesus IFN- γ , 59%-64% with bovine, canine, equine, feline, and porcine IFN- γ , and 37%-43% with cotton rat, mouse, and rat IFN- γ . IFN- γ dimers bind to IFN- γ RI (α subunits) which then interact with IFN- γ RII (β subunits) to form the functional receptor complex of two α and two β subunits. Inclusion of IFN- γ RII increases the binding affinity for ligand and the efficiency of signal transduction (4, 5). IFN- γ is produced by a variety of immune cells under inflammatory conditions, notably by T cells and NK cells (6). It plays a key role in host defense by promoting the development and activation of Th1 cells, chemoattraction and activation of monocytes and macrophages, up-regulation of antigen presentation molecules, and immunoglobulin class switching in B cells. It also exhibits antiviral, antiproliferative, and apoptotic effects (6, 7). In addition, IFN- γ functions as an anti-inflammatory mediator by promoting the development of regulatory T cells and inhibiting Th17 cell differentiation (8, 9). The pleiotropic effects of IFN- γ contribute to the development of multiple aspects of atherosclerosis (7).

References:

1. Billiau, A. and P. Matthys (2009) Cytokine Growth Factor Rev. **20**:97.
2. Pestka, S. *et al.* (2004) Immunol. Rev. **202**:8.
3. Gray, P.W. and D.V. Goeddel (1982) Nature **298**:859.
4. Marsters, S.A. *et al.* (1995) Proc. Natl. Acad. Sci. **92**:5401.
5. Krause, C.D. *et al.* (2000) J. Biol. Chem. **275**:22995.
6. Schroder, K. *et al.* (2004) J. Leukoc. Biol. **75**:163.
7. McLaren, J.E. and D.P. Ramji (2009) Cytokine Growth Factor Rev. **20**:125.
8. Muhl, H. and J. Pfeilschifter (2003) Int. Immunopharmacol. **3**:1247.
9. Kelchtermans, H. *et al.* (2008) Trends Immunol. **29**:479.