

DESCRIPTION

Source *Spodoptera frugiperda*, Sf 21 (baculovirus)-derived
Thr27-Pro185
Accession # AAA17999.1

N-terminal Sequence Analysis Thr27

Predicted Molecular Mass 17.5 kDa

SPECIFICATIONS

SDS-PAGE 17-30 kDa, reducing conditions

Activity Measured in a cell proliferation assay using BaF3 mouse pro-B cells transfected with mouse Flt-3. The ED₅₀ for this effect is typically 0.2-1 ng/mL. The specific activity of Recombinant Human Flt-3 Ligand is approximately 1.2 x 10³ U/μg, which is calibrated against recombinant human Flt-3 Ligand WHO Standard (NIBSC code: 96/532).

Endotoxin Level <0.10 EU per 1 μg of the protein by the LAL method.

Purity >97%, by SDS-PAGE visualized with Silver Staining and quantitative densitometry by Coomassie® Blue Staining.

Formulation Lyophilized from a 0.2 μm filtered solution in Acetonitrile and TFA. See Certificate of Analysis for details.

PREPARATION AND STORAGE



Reconstitution Reconstitute 5 μg vials at 50 μg/mL in sterile PBS. Reconstitute 25 μg or larger vials at 100 μg/mL in sterile PBS.

Shipping The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

- 12 months from date of receipt, -20 to -70 °C as supplied.
- 1 month, 2 to 8 °C under sterile conditions after reconstitution.
- 3 months, -20 to -70 °C under sterile conditions after reconstitution.

DATA



BACKGROUND

Flt-3 Ligand, also known as FL, is an α -helical cytokine that promotes the differentiation of multiple hematopoietic cell lineages (1 - 3). Mature human Flt-3 Ligand consists of a 158 amino acid (aa) extracellular domain (ECD) with a cytokine-like domain and a juxtamembrane tether region, a 21 aa transmembrane segment, and a 30 aa cytoplasmic tail (4 - 7). Within the ECD, human Flt-3 Ligand shares 71% and 65% aa sequence identity with mouse and rat Flt-3 Ligand, respectively. Human and mouse Flt-3 Ligand show cross-species activity (4 - 6). Flt-3 Ligand is expressed as a noncovalently-linked dimer by T cells and bone marrow and thymic fibroblasts (1, 8). Each 36 kDa chain carries approximately 12 kDa of N- and O-linked carbohydrates (8). Alternate splicing and proteolytic cleavage of the transmembrane form can generate a soluble 30 kDa fragment that includes the cytokine domain (4, 8). Alternate splicing of human Flt-3 Ligand also generates membrane-associated isoforms that contain either a truncated cytoplasmic tail or an 85 aa substitution following the cytokine domain (4, 5, 8). Both transmembrane and soluble Flt-3 Ligand signal through the tyrosine kinase receptor Flt-3/Flk-2 (3, 4, 6, 7). Flt-3 Ligand induces the expansion of monocytes and immature dendritic cells as well as early B cell lineage differentiation (2, 9). It synergizes with IL-3, GM-CSF, and SCF to promote the mobilization and myeloid differentiation of hematopoietic stem cells (4 - 6). It cooperates with IL-2, -6, -7, and -15 to induce NK cell development and with IL-3, -7, and -11 to induce terminal B cell maturation (1, 10). Animal studies also show Flt-3 Ligand to reduce the severity of experimentally induced allergic inflammation (11).

References:

1. Wodnar-Filipowicz, A. (2003) *News Physiol. Sci.* **18**:247.
2. Dong, J. *et al.* (2002) *Cancer Biol. Ther.* **1**:486.
3. Gilliland, D.G. and J.D. Griffin (2002) *Blood* **100**:1532.
4. Hannum, C. *et al.* (1994) *Nature* **368**:643.
5. Lyman, S.D. *et al.* (1994) *Blood* **83**:2795.
6. Lyman, S.D. *et al.* (1993) *Cell* **75**:1157.
7. Savvides, S.N. *et al.* (2000) *Nat. Struct. Biol.* **7**:486.
8. McClanahan, T. *et al.* (1996) *Blood* **88**:3371.
9. Diener, K.R. *et al.* (2008) *Exp. Hematol.* **36**:51.
10. Farag, S.S. and M.A. Caligiuri (2006) *Blood Rev.* **20**:123.
11. Edwan, J.H. *et al.* (2004) *J. Immunol.* **172**:5016.