

**DESCRIPTION**

<b>Source</b>	<i>E. coli</i> -derived	
	Human BMP-2 (Ala284 – Arg396) Accession # NP_001191	
	Human BMP-7 (Ser293 – His431) Accession # NP_001710	
	N-terminus	C-terminus

<b>N-terminal Sequence Analysis</b>	Ala284 & Ser293
<b>Structure / Form</b>	Disulfide-linked heterodimer
<b>Predicted Molecular Mass</b>	12.9 kDa (BMP-2), 15.8 kDa (BMP-7)

**SPECIFICATIONS**

<b>Activity</b>	Measured by its ability to induce alkaline phosphatase production by ATDC5 mouse chondrogenic cells. Nakamura, K. <i>et al.</i> (1999) <i>Exp. Cell Res.</i> <b>250</b> :351. The ED <sub>50</sub> for this effect is 10-40 ng/mL.
<b>Endotoxin Level</b>	<0.10 EU per 1 µg of the protein by the LAL method.
<b>Purity</b>	>97%, by SDS-PAGE under reducing conditions and visualized by silver stain.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in Acetonitrile and TFA. See Certificate of Analysis for details.

**PREPARATION AND STORAGE**

<b>Reconstitution</b>	Reconstitute at 100 µg/mL in sterile 4 mM HCl.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>● 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>● 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>● 3 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

**BACKGROUND**

Human BMP-7, also known as osteogenic protein 1 (OP-1), and BMP-2 are members of the BMP subgroup of the TGF-β superfamily and signal through heterodimeric complexes composed of type I and type II BMP receptors. BMP-2 and BMP-7 influence a variety of morphogenic processes, particularly during skeletal and renal development (1 - 3). The human BMP-2 cDNA encodes a 396 amino acid (aa) precursor that contains a 23 aa signal sequence, a 259 propeptide, and a 114 aa mature protein (4). The human BMP-7 cDNA encodes a 431 aa precursor that contains a 29 aa signal sequence, a 263 aa propeptide, and a 139 aa mature protein (5). BMP propeptides are removed by proteolysis, enabling mature BMPs to form active disulfide linked homodimers or heterodimers (1). Human and mouse BMP-2 and BMP-7 are 100% and 98% identical, respectively, at the amino acid level. Human BMP-2 shares 85% aa sequence identity with human BMP-4 and less than 51% aa sequence identity with other BMPs. Human BMP-7 shares approximately 60% - 70% aa sequence identity with BMP-5, -6, and -8, and less than 50% aa sequence identity with other BMPs. BMP-2 and BMP-7 are co-expressed in some embryonic tissues (6, 7) and associate into a functional 38 kDa osteogenic dimer (8). In *in vitro* osteoblast differentiation assays and *in vivo* bone formation models, a BMP-2/BMP-7 heterodimer is significantly more potent than either homodimer (9 - 12). Considering that BMP-2 preferentially binds BMPRIA/ALK-3 and BMPRII/ALK-6, while BMP-7 is selective for ALK-2, the observed increase in heterodimer activity may be due to the triggering of additional receptor subtypes.

**References:**

1. Chen, D. *et al.* (2004) *Growth Factors* **22**:233.
2. Miyazono, K. *et al.* (2005) *Cytokine Growth Factor Rev.* **16**:251.
3. Simic, P. and S. Vukicevic (2005) *Cytokine Growth Factor Rev.* **16**:299.
4. Wozney, J. *et al.* (1988) *Science* **242**:1528.
5. Gregory, K.E. *et al.* (2005) *J. Biol. Chem.* **280**:27970.
6. Lyons, K.M. *et al.* (1995) *Mech. Dev.* **50**:71.
7. Knosp, W.M. *et al.* (2004) *Development* **131**:4581.
8. Sampath, T.K. *et al.* (1990) *J. Biol. Chem.* **265**:13198.
9. Hazama, M. *et al.* (1995) *Biochem. Biophys. Res. Commun.* **209**:859.
10. Zhu, W. *et al.* (2004) *J. Bone Miner. Res.* **19**:2021.
11. Zhao, M. *et al.* (2005) *J. Cell. Biochem.* **95**:1.
12. Israel, D.I. *et al.* (1996) *Growth Factors* **13**:291.