

DESCRIPTION

Source	Chinese Hamster Ovary cell line, CHO-derived BMP-2 protein Gln283-Arg396 Accession # P12643
Structure / Form	Disulfide-linked homodimer
Predicted Molecular Mass	13 kDa (monomer)


SPECIFICATIONS

SDS-PAGE	15-16 kDa, reducing conditions
Activity	Measured by its ability to induce alkaline phosphatase production by ATDC5 mouse chondrogenic cells. Binnerts, M.E. <i>et al.</i> (2004) <i>Biochem. Biophys. Res. Commun.</i> 315 (2):272. The ED ₅₀ for this effect is 40-200 ng/mL. The specific activity of Recombinant Human BMP-2 is approximately 781 U/μg, which is calibrated against human BMP-2 WHO International Standard (NIBSC code: 93/574). Specific activity is for reference purposes only and is not routinely tested.
Endotoxin Level	<0.10 EU per 1 μg of the protein by the LAL method.
Purity	>95%, by SDS-PAGE under reducing conditions and visualized by silver stain.
Formulation	Lyophilized from a 0.2 μm filtered solution in Glycine, Sucrose, Tween® 80 and Glutamic Acid with BSA as a carrier protein. *1 mg pack size (01M) is supplied as a 0.2 μm filtered solution in Glycine, Sucrose, Tween® 80 and Glutamic Acid with BSA as a carrier protein. See Certificate of Analysis for details.

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 100-200 μg/mL in sterile 4 mM HCl containing at least 0.1% human or bovine serum albumin.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> • 12 months from date of receipt, -20 to -70 °C as supplied. • 1 month, 2 to 8 °C under sterile conditions after reconstitution. • 3 months, -20 to -70 °C under sterile conditions after reconstitution.

DATA



BACKGROUND

Bone morphogenetic protein 2 (BMP-2) is a member of the BMP subgroup of the TGF- β superfamily. It plays a dominant role in embryonic dorsal-ventral patterning, organogenesis, limb bud formation, and bone formation and regeneration (1, 2). Human BMP-2 is synthesized as a 396 amino acid (aa) preproprotein that contains a 23 aa signal sequence, a 259 aa prosegment, and a 114 aa mature region (3). Proteolytic removal of the propeptide enables mature BMP-2 to form active disulfide linked homodimers and heterodimers with BMP-7 (2). Mature monomeric BMP-2 is an 18 kDa glycosylated peptide with seven conserved cysteines that form a cystine knot structure (4). Mature human BMP-2 shares 100% aa sequence identity with mouse and rat BMP-2. It shares 85% aa sequence identity with human BMP-4 and less than 51% with other BMPs. BMP-2 signals through heterodimeric complexes composed of a type I receptor (Activin RI, BMPRI-IA, or BMPRI-IB) and a type II receptor (BMPRII or ActivinRIIB) (2, 5). BMP-2 induces chondrocyte proliferation, endochondral bone formation, longitudinal bone growth, and bone and cartilage repair (6, 7). It induces ectopic bone formation or calcification by promoting osteogenic and chondrogenic differentiation in mesenchymal cells, stem cells, and vascular smooth muscle cells (2, 8-10). BMP-2/BMP-7 heterodimers are significantly more potent than BMP-2 homodimers at inducing bone formation *in vivo* (11). BMP-2 also promotes the maintenance and repair of colonic epithelium, suppresses neuronal dopamine synthesis and release, induces apoptosis in medulloblastoma cells, and is required for cardiac contractility (12-15).

References:

1. Kishigami, S. and Y. Mishina (2005) Cytokine Growth Factor Rev. **16**:265.
2. Chen, D. *et al.* (2004) Growth Factors **22**:233.
3. Wozney, J. *et al.* (1988) Science **242**:1528.
4. Sun, P.D. and D.R. Davies (1995) Annu. Rev. Biophys. Biomol. Struct. **24**:269.
5. Sebald, W. *et al.* (2004) Biol. Chem. **385**:697.
6. De Luca, F. *et al.* (2001) Endocrinology **142**:430.
7. Davidson, E.N.B., *et al.* (2007) Arthritis Res. Ther. **9**:R102.
8. Ryoo, H.-M. *et al.* (2006) Gene **366**:51.
9. Kramer, J. *et al.* (2000) Mech. Dev. **92**:193.
10. Li, X. *et al.* (2008) Atherosclerosis January 5 epub.
11. Zhu, W. *et al.* (2004) J. Bone Miner. Res. **19**:2021.
12. Peiris, D. *et al.* (2007) Am. J. Physiol. Gastrointest. Liver Physiol. **292**:G753.
13. Kano, Y. *et al.* (2005) Endocrinology **146**:5332.
14. Hallahan, A.R. *et al.* (2003) Nat. Med. **9**:1033.
15. Wang, Y.-X. *et al.* (2007) Cardiovasc. Res. **74**:290.