Recombinant Human Active NEK2

Certificate of Analysis Catalog Number: 3706-KS Lot Number: 1556939

Specifications and Use

Source	 Recombinant full-length human NEK2 was expressed by baculovirus in <i>Sf</i> 9 insect cells using an N-terminal GST tag. The gene accession number is NM_002497.
Molecular Mass	• The approximate molecular weight is 76 kDa (see Figure 1 below).
Purity	 The purity was determined to be > 90% by densitometry (see Figure 1 below).
Formulation	 Supplied in 50 mM Tris-HCl, pH 7.5, 150 mM NaCl, 0.25 mM DTT, 0.1 mM EGTA, 0.1 mM EDTA, 0.1 mM PMSF, 25% glycerol.
Size	 ♦ 10 μg.
Concentration	 ♦ 0.1 μg/μL.
Activity	 The specific activity of NEK2 was determined to be 270 nmol/min/mg using a myelin basic protein (MBP) substrate (see activity assay protocol).
Storage	 This product is stable at ≤ -70 °C for up to 1 year from the date of receipt. For optimal storage, aliquot into smaller quantities after centrifugation and store at recommended temperature. Avoid repeated freeze-thaw cycles.

NEK2

NEK2 is closely related in its catalytic domain to the Serine/Threonine protein kinase NIMA of *Aspergillus nidulans* that is required for entry into mitosis and may function in parallel to the universal mitotic inducer p34cdc2. Like NIMA, the NEK2 protein is almost undetectable during G1 but accumulates progressively throughout S, reaching maximal levels in late G2 (1). NEK2 is shown to be expressed most abundantly in the testis of the adult tissues examined being localized to the nucleus (2).

References

- 1. Schultz, S.J. et al. (1994) Cell Growth Differ. 5:625.
- 2. Fry, A.M. *et al.* (1995) J. Biol. Chem. **270**:12899.







Figure 2: Enzymatic assay results. The specific activity of NEK2 was determined to be 270 nmol/min/mg as per activity assay protocol (on reverse).

FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

R&D Systems, Inc. 1-800-343-7475

Activity Assay Protocol

Solutions Required

- Active Kinase Active NEK2 (0.1 μg/μL) diluted with Kinase Dilution Buffer and assayed as outlined in Figure 2.
 Note: These are suggested working dilutions. Optimal dilutions should be determined by each laboratory for each application.
- Kinase Assay Buffer I, pH 7.2 25 mM MOPS, 12.5 mM β-glycerolphosphate, 25 mM MgCl₂, 5 mM EGTA, 2 mM EDTA. Add 0.25 mM DTT to the Kinase Assay Buffer prior to use.
- Kinase Dilution Buffer, pH 7.2 Kinase Assay Buffer I diluted at a 1:4 ratio (5-fold dilution) with 50 ng/µL BSA solution.
- 10 mM ATP Stock Solution Prepare the ATP Stock Solution by dissolving 55 mg of ATP in 10 mL of Kinase Assay Buffer I.
- [³³P]-ATP Assay Cocktail Prepare 250 μM [³³P]-ATP Assay Cocktail in a designated radioactive work area by combining 150 μL of 10 mM ATP Stock Solution, 100 μL of [³³P]-ATP (1 mCi/100 μL), and 5.75 mL of Kinase Assay Buffer I.
- Substrate Myelin basic protein (MBP) substrate diluted in distilled or deionized water to a final concentration of 1 mg/mL.

Assay Procedure

- 1. Thaw the [³³P]-ATP Assay Cocktail in a shielded container in a designated radioactive work area.
- 2. Thaw the Active NEK2, Kinase Assay Buffer I, Substrate, and Kinase Dilution Buffer on ice.
- 3. In a pre-cooled microfuge tube, add the following reaction components bringing the initial reaction volume up to 20 µL.

Reaction Component	Amount
Diluted Active NEK2	10 μL
Substrate (1 mg/mL Stock Solution)	5 μL
Distilled H ₂ O (2-8 °C)	5 μL

- 4. Set up the blank control as outlined in step 3, excluding the addition of the substrate. Replace the substrate with an equal volume of distilled or deionized water.
- 5. Initiate the reaction by the addition of 5 μL [³³P]-ATP Assay Cocktail, bringing the final volume up to 25 μL. Incubate the mixture in a water bath at 30 °C for 15 minutes.
- 6. After the 15 minute incubation period, terminate the reaction by spotting 20 μL of the reaction mixture onto individual pre-cut strips of phosphocellulose P81 paper.
- 7. Air dry the pre-cut P81 strip and sequentially wash in a 1% phosphoric acid solution (dilute 10 mL of phosphoric acid and make a 1 liter solution with distilled or deionized water) with constant gentle stirring. It is recommended that the strips be washed a total of three times for approximately 10 minutes each.
- 8. Count the radioactivity on the P81 paper in the presence of scintillation fluid in a scintillation counter.
- 9. Determine the corrected cpm by removing the blank control value (see step 4) for each sample and calculate the kinase specific activity as outlined below.

Calculation of [³³P]-ATP Specific Activity (SA) (cpm/pmol)

Specific Activity (SA) = cpm for 5 µL [³³P]-ATP/pmole of ATP (in 5 µL of a 250 µM ATP stock solution, *i.e.* 1250 pmoles)

Calculation of Kinase Specific Activity (SA) (pmol/minutes/µg or nmol/minutes/mg)

Corrected cpm from reaction / [(SA of ³³P-ATP in cpm/pmol) x (Reaction time in minutes) x (Enzyme amount in μ g or mg)] x [(Reaction volume) / (Spot Volume)]

Diane Wotta

pliane R. Wola

Quality & Regulatory Affairs