

DESCRIPTION

Source *E. coli*-derived
Val118-Ser269, with an N-terminal Met
Accession # NP_032387

N-terminal Sequence Analysis Met

Predicted Molecular Mass 17 kDa

SPECIFICATIONS

Activity Measured in a cell proliferation assay using D10.G4.1 mouse helper T cells. Symons, J.A. *et al.* (1987) in *Lymphokines and Interferons*, a Practical Approach. Clemens, M.J. *et al.* (eds): IRL Press. 272.
The ED₅₀ for this effect is typically 2-10 μ g/mL.

Endotoxin Level <0.01 EU per 1 μ g of the protein by the LAL method.

Purity >97%, by SDS-PAGE under reducing conditions and visualized by silver stain.

Formulation Lyophilized from a 0.2 μ m filtered solution in PBS with BSA as a carrier protein. See Certificate of Analysis for details.

PREPARATION AND STORAGE



Reconstitution Reconstitute at 100 μ g/mL in sterile PBS containing at least 0.1% human or bovine serum albumin.

Shipping The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

- 12 months from date of receipt, -20 to -70 °C as supplied.
- 1 month, 2 to 8 °C under sterile conditions after reconstitution.
- 3 months, -20 to -70 °C under sterile conditions after reconstitution.

DATA



BACKGROUND

IL-1 is a name that designates two pleiotropic cytokines, IL-1 α (IL-1F1) and IL-1 β (IL-1F2), which are the products of distinct genes. IL-1 α and IL-1 β are structurally related polypeptides that share approximately 17% amino acid (aa) identity in mouse. Both proteins are produced by a wide variety of cells in response to inflammatory agents, infections, or microbial endotoxins. While IL-1 α and IL-1 β are regulated independently, they bind to the same receptor and exert identical biological effects. IL-1 RI binds directly to IL-1 α or IL-1 β and then associates with IL-1 R accessory protein (IL-1 R3/IL-1 RAcP) to form a high-affinity receptor complex that is competent for signal transduction. IL-1 RII has high affinity for IL-1 β but functions as a decoy receptor and negative regulator of IL-1 β activity. IL-1ra functions as a competitive antagonist by preventing IL-1 α and IL-1 β from interacting with IL-1 RI (1-4). The mouse IL-1 β cDNA encodes a 269 aa precursor. A 117 aa propeptide is cleaved intracellularly by the cysteine protease IL-1 β -converting enzyme (Caspase-1/ICE) to generate the active cytokine (5, 6). The 17 kDa mature mouse IL-1 β shares 90% aa sequence identity with cotton rat and rat and 65%-78% identity with canine, equine, feline, human, porcine, and rhesus IL-1 β .

References:

1. Allan, S.M. *et al.* (2005) *Nat. Rev. Immunol.* **5**:629.
2. Boraschi, D. and A. Tagliabue (2006) *Vitam. Horm.* **74**:229.
3. Kornman, K.S. (2006) *Am. J. Clin. Nutr.* **83**:475S.
4. Isoda, K. and F. Ohsuzu (2006) *J. Atheroscler. Thromb.* **13**:21.
5. Gray, P.W. *et al.* (1986) *J. Immunol.* **137**:3644.
6. Martinon, F. and J. Tschopp (2007) *Cell Death Differ.* **14**:10.