Recombinant Mouse IL-4

Catalog Number: 404-ML

DESCRIPTION

Source	*E. coli*-derived His23-Ser140, with an N-terminal Met Accession # P07750
N-terminal Sequence Analysis	His23
Predicted Molecular Mass	14 kDa

SPECIFICATIONS

Activity	Measured in a cell proliferation assay using HT-2 mouse T cells. The ED_{50} for this effect is typically 0.3-1.5 ng/mL.
Endotoxin Level	<0.10 EU per 1 µg of the protein by the LAL method.
Purity	>97%, by SDS-PAGE visualized with Silver Staining and quantitative densitometry by Coomassie® Blue Staining.
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with BSA as a carrier protein. See Certificate of Analysis for details.

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 100 µg/mL in sterile PBS containing at least 0.1% human or bovine serum albumin.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	**Use a manual defrost freezer and avoid repeated freeze-thaw cycles.** ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 3 months, -20 to -70 °C under sterile conditions after reconstitution.

DATA

SDS-PAGE



1 µg/lane of Recombinant Mouse IL-4 was resolved with SDS-PAGE under reducing (R) conditions and visualized by silver staining, showing a single band at 12 kDa.

Bioactivity



Recombinant Mouse IL-4 (Catalog # 404-ML) stimulates cell proliferation of the HT-2 mouse T cell line. The ED_{50} for this effect is typically 0.3-1.5 ng/mL.

bio-techne
bio-techne.com
Global info@bio-techne.com techsupport@bio-techne.com
North America TEL 800 343 7475 Europe | Middle East | Africa TEL +44 (0)1235 529449
China info.cn@bio-techne.com TEL +86 (21) 52380373
Rest of World bio-techne.com/find-us/distributors TEL +1 612 379 2956

BACKGROUND

Interleukin-4 (IL-4), also known as B cell-stimulatory factor-1, is a monomeric, approximately 13 kDa-18 kDa Th2 cytokine that shows pleiotropic effects during immune responses (1-4). It is a glycosylated polypeptide that contains three intrachain disulfide bridges and adopts a bundled four α-helix structure (5). Mouse IL-4 is synthesized with a 24 aa signal sequence. Mature mouse IL-4 shares 39%, 39%, and 59% aa sequence identity with bovine, human, and rat IL-4, respectively. Human, mouse, and rat IL-4 are species-specific in their activities (6-8). IL-4 exerts its effects through two receptor complexes (9, 10). The type I receptor, which is expressed on hematopoietic cells, is a heterodimer of the ligand binding IL-4 Rα and the common γ chain (a shared subunit of the receptors for IL-2, -7, -9, -15, and -21). The type II receptor on nonhematopoietic cells consists of IL-4 Rα and IL-13 Rα1. The type II receptor also transduces IL-13 mediated signals. IL-4 is primarily expressed by Th2-biased $CD4^+$ T cells, mast cells, basophils, and eosinophils (1, 2). It promotes cell proliferation, survival, and immunoglobulin class switch to IgG1 and IgE in mouse B cells, acquisition of the Th2 phenotype by naïve $CD4^+$ T cells, priming and chemotaxis of mast cells, eosinophils, and basophils, and the proliferation and activation of epithelial cells (11 - 14). IL-4 plays a dominant role in the development of allergic inflammation and asthma (13, 15).

References:

1. Benczik, M. and S.L. Gaffen (2004) Immunol. Invest. **33**:109.
2. Chomarat, P. and J. Banchereau (1998) Int. Rev. Immunol. **17**:1.
3. Lee, F. *et al.* (1986) Proc. Natl. Acad. Sci. **83**:2061.
4. Noma, Y. *et al.* (1986) Nature **319**:640.
5. Redfield, C. *et al.* (1991) Biochemistry **30**:11029.
6. Ramirez, F. *et al.* (1988) J. Immunol. Meth. **221**:141.
7. Leitenberg, D. and T.L. Feldbush (1988) Cell. Immunol. **111**:451.
8. Mosman, T.R. *et al.* (1987) J. Immunol. **138**:1813.
9. Mueller, T.D. *et al.* (2002) Biochim. Biophys. Acta **1592**:237.
10. Nelms, K. *et al.* (1999) Annu. Rev. Immunol. **17**:701.
11. Paludan, S.R. (1998) Scand. J. Immunol. **48**:459.
12. Corthay, A. (2006) Scand. J. Immunol. **64**:93.
13. Ryan, J.J. *et al.* (2007) Crit. Rev. Immunol. **27**:15.
14. Grone, A. (2002) Vet. Immunol. Immunopathol. **88**:1.
15. Rosenberg, H.F. *et al.* (2007) J. Allergy Clin. Immunol. **119**:1303.

bio-techne
bio-techne.com
Global info@bio-techne.com techsupport@bio-techne.com
North America TEL 800 343 7475 Europe | Middle East | Africa TEL +44 (0)1235 529449
China info.cn@bio-techne.com TEL +86 (21) 52380373
Rest of World bio-techne.com/find-us/distributors TEL +1 612 379 2956