

DESCRIPTION

Source *E. coli*-derived
His23-Ser140, with an N-terminal Met
Accession # P07750

N-terminal Sequence Analysis His23

Predicted Molecular Mass 14 kDa

SPECIFICATIONS

Activity Measured in a cell proliferation assay using HT-2 mouse T cells.
The ED₅₀ for this effect is 0.3-1.5 ng/mL.

Endotoxin Level <0.10 EU per 1 µg of the protein by the LAL method.

Purity >97%, by SDS-PAGE visualized with Silver Staining and quantitative densitometry by Coomassie® Blue Staining.

Formulation Lyophilized from a 0.2 µm filtered solution in PBS. See Certificate of Analysis for details.

PREPARATION AND STORAGE

Reconstitution Reconstitute at 100 µg/mL in sterile PBS.


Shipping The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

- 12 months from date of receipt, -20 to -70 °C as supplied.
- 1 month, 2 to 8 °C under sterile conditions after reconstitution.
- 3 months, -20 to -70 °C under sterile conditions after reconstitution.


DATA

Bioactivity



Recombinant Mouse IL-4 (Catalog # 404-ML/CF) stimulates cell proliferation of the HT-2 mouse T cell line. The ED₅₀ for this effect is 0.3-1.5 ng/mL.

SDS-PAGE



1 µg/lane of Recombinant Mouse IL-4 was resolved with SDS-PAGE under reducing (R) conditions and visualized by silver staining, showing a single band at 12 kDa.

BACKGROUND

Interleukin-4 (IL-4), also known as B cell-stimulatory factor-1, is a monomeric, approximately 13 kDa-18 kDa Th2 cytokine that shows pleiotropic effects during immune responses (1-4). It is a glycosylated polypeptide that contains three intrachain disulfide bridges and adopts a bundled four α -helix structure (5). Mouse IL-4 is synthesized with a 24 aa signal sequence. Mature mouse IL-4 shares 39%, 39%, and 59% aa sequence identity with bovine, human, and rat IL-4, respectively. Human, mouse, and rat IL-4 are species-specific in their activities (6-8). IL-4 exerts its effects through two receptor complexes (9, 10). The type I receptor, which is expressed on hematopoietic cells, is a heterodimer of the ligand binding IL-4 R α and the common γ chain (a shared subunit of the receptors for IL-2, -7, -9, -15, and -21). The type II receptor on nonhematopoietic cells consists of IL-4 R α and IL-13 R α 1. The type II receptor also transduces IL-13 mediated signals. IL-4 is primarily expressed by Th2-biased CD4⁺ T cells, mast cells, basophils, and eosinophils (1, 2). It promotes cell proliferation, survival, and immunoglobulin class switch to IgG1 and IgE in mouse B cells, acquisition of the Th2 phenotype by naïve CD4⁺ T cells, priming and chemotaxis of mast cells, eosinophils, and basophils, and the proliferation and activation of epithelial cells (11 - 14). IL-4 plays a dominant role in the development of allergic inflammation and asthma (13, 15).

References:

1. Benczik, M. and S.L. Gaffen (2004) *Immunol. Invest.* **33**:109.
2. Chomarat, P. and J. Banchereau (1998) *Int. Rev. Immunol.* **17**:1.
3. Lee, F. *et al.* (1986) *Proc. Natl. Acad. Sci.* **83**:2061.
4. Noma, Y. *et al.* (1986) *Nature* **319**:640.
5. Redfield, C. *et al.* (1991) *Biochemistry* **30**:11029.
6. Ramirez, F. *et al.* (1988) *J. Immunol. Meth.* **221**:141.
7. Leitenberg, D. and T.L. Feldbush (1988) *Cell. Immunol.* **111**:451.
8. Mosman, T.R. *et al.* (1987) *J. Immunol.* **138**:1813.
9. Mueller, T.D. *et al.* (2002) *Biochim. Biophys. Acta* **1592**:237.
10. Nelms, K. *et al.* (1999) *Annu. Rev. Immunol.* **17**:701.
11. Paludan, S.R. (1998) *Scand. J. Immunol.* **48**:459.
12. Corthay, A. (2006) *Scand. J. Immunol.* **64**:93.
13. Ryan, J.J. *et al.* (2007) *Crit. Rev. Immunol.* **27**:15.
14. Grone, A. (2002) *Vet. Immunol. Immunopathol.* **88**:1.
15. Rosenberg, H.F. *et al.* (2007) *J. Allergy Clin. Immunol.* **119**:1303.