

DESCRIPTION

Source	<i>E. coli</i> -derived mouse M-CSF protein Lys33-Glu262, with an N-terminal Met Accession # Q3U4F9
N-terminal Sequence Analysis	Met
Structure / Form	Disulfide-linked homodimer
Predicted Molecular Mass	26 kDa (monomer)


SPECIFICATIONS

SDS-PAGE	29 kDa, reducing conditions
Activity	Measured in a cell proliferation assay using M-NFS-60 mouse myelogenous leukemia lymphoblast cells. Nakoinz, I. <i>et al.</i> (1990) J. Immunol. 145:860. The ED ₅₀ for this effect is 0.5-3 ng/mL.
Endotoxin Level	<1.0 EU per 1 µg of the protein by the LAL method.
Purity	>97%, by SDS-PAGE under reducing conditions and visualized by silver stain.
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with BSA as a carrier protein. See Certificate of Analysis for details.

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 100 µg/mL in sterile PBS containing at least 0.1% human or bovine serum albumin.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> • 12 months from date of receipt, -20 to -70 °C as supplied. • 1 month, 2 to 8 °C under sterile conditions after reconstitution. • 3 months, -20 to -70 °C under sterile conditions after reconstitution.

DATA



BACKGROUND

M-CSF, also known as CSF-1, is a four- α -helical-bundle cytokine that is the primary regulator of macrophage survival, proliferation and differentiation (1-3). M-CSF protein is also essential for the survival and proliferation of osteoclast progenitors (1, 4). M-CSF also primes and enhances macrophage killing of tumor cells and microorganisms, regulates the release of cytokines and other inflammatory modulators from macrophages, and stimulates pinocytosis (2, 3). M-CSF increases during pregnancy to support implantation and growth of the decidua and placenta (5). Sources of M-CSF include fibroblasts, activated macrophages, endometrial secretory epithelium, bone marrow stromal cells and activated endothelial cells (1-5). The M-CSF receptor (*c-fms*) transduces its pleiotropic effects and mediates its endocytosis. M-CSF mRNAs of various sizes occur (3-9). Full length mouse M-CSF transcripts encode a 520 amino acid (aa) type I transmembrane (TM) protein with a 462 aa extracellular region, a 21 aa TM domain, and a 37 aa cytoplasmic tail that forms a 140 kDa covalent dimer. Differential processing produces two proteolytically cleaved, secreted dimers. One is an N- and O- glycosylated 86 kDa dimer, while the other is modified by both glycosylation and chondroitin-sulfate proteoglycan (PG) to generate a 200 kDa subunit. Although PG-modified M-CSF protein can circulate, it may be immobilized by attachment to type V collagen (8). Shorter transcripts encode M-CSF that lacks cleavage and PG sites and produces an N-glycosylated 68 kDa TM dimer and a slowly produced 44 kDa secreted dimer (7). Although forms may vary in activity and half-life, all contain the N-terminal 150 aa portion that is necessary and sufficient for interaction with the M-CSF receptor (10, 11). The first 229 aa of mature mouse M-CSF shares 87%, 83%, 82% and 81% aa identity with corresponding regions of rat, dog, cow and human M-CSF, respectively (12, 13). Human M-CSF is active in the mouse, but mouse M-CSF is reported to be species-specific.

References:

1. Pixley, F.J. and E.R. Stanley (2004) Trends Cell Biol. **14**:628.
2. Chitu, V. and E.R. Stanley (2006) Curr. Opin. Immunol. **18**:39.
3. Fixe, P. and V. Praloran (1997) Eur. Cytokine Netw. **8**:125.
4. Ryan, G.R. *et al.* (2001) Blood **98**:74.
5. Makrigiannakis, A. *et al.* (2006) Trends Endocrinol. Metab. **17**:178.
6. Nandi, S. *et al.* (2006) Blood **107**:786.
7. Rettenmier, C.W. and M.F. Roussel (1988) Mol. Cell Biol. **8**:5026.
8. Suzu, S. *et al.* (1992) J. Biol. Chem. **267**:16812.
9. Manos, M.M. (1988) Mol. Cell. Biol. **8**:5035.
10. Koths, K. (1997) Mol. Reprod. Dev. **46**:31.
11. Jang, M-H. *et al.* (2006) J. Immunol. **177**:4055.
12. DeLamarter, J.F. *et al.* (1987) Nucleic Acids Res. **15**:2389.
13. Ladner, M.B. *et al.* (1988) Proc. Natl. Acad. Sci. USA **85**:6706.