

DESCRIPTION

Source *E. coli*-derived mouse Sonic Hedgehog/Shh protein
Cys25-Gly198 (Cys25Ile-Ile), with an N-terminal Met
Accession # Q62226

N-terminal Sequence Analysis Met

Predicted Molecular Mass 19.8 kDa

SPECIFICATIONS

Activity Measured by its ability to induce alkaline phosphatase production by C3H10T1/2 mouse embryonic fibroblast cells. Nakamura, T. *et al.* (1997) Biochem. Biophys. Res. Commun. **237**:465.
The ED₅₀ for this effect is 0.05-0.25 µg/mL.

Endotoxin Level <0.10 EU per 1 µg of the protein by the LAL method.

Purity >97%, by SDS-PAGE under reducing conditions and visualized by silver stain.

Formulation Lyophilized from a 0.2 µm filtered solution in NaH₂PO₄, NaCl and DTT. See Certificate of Analysis for details.

PREPARATION AND STORAGE

Reconstitution Reconstitute at 100-200 µg/mL in sterile PBS.


Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

- 12 months from date of receipt, -20 to -70 °C as supplied.
- 1 month, 2 to 8 °C under sterile conditions after reconstitution.
- 3 months, -20 to -70 °C under sterile conditions after reconstitution.


DATA

Bioactivity



Recombinant Mouse Sonic Hedgehog/Shh (C25II) N-Terminus Bioactivity
Recombinant Mouse Sonic Hedgehog/Shh (C25II), N-Terminus (Catalog # 464-SH/CF) induces alkaline phosphatase production by the C3H10T1/2 mouse embryonic fibroblast cell line. The activity is more than 30-fold greater than the top competitor's Sonic Hedgehog.

SDS-PAGE



Recombinant Mouse Sonic Hedgehog/Shh (C25II) N-Terminus SDS-PAGE 1 µg/lane of Recombinant Mouse Sonic Hedgehog/Shh (C25II), N-Terminus was resolved with SDS-PAGE under reducing (R) conditions and visualized by silver staining, showing a single band at 22 kDa.

BACKGROUND

Sonic Hedgehog (Shh) is expressed in embryonic tissues that are critical for the patterning of the developing central nervous system, somite, and limb. It is also involved in whisker, hair, foregut, tooth, and bone development. Shh regulates neural and hematopoietic stem cell fate and is important for thymocyte differentiation and proliferation as well as T cell determination. In adult tissue Shh is associated with cancer development and tissue remodeling following injury (1-3). Mouse Shh encodes a 437 amino acid (aa) precursor protein that is autocatalytically processed to yield a non-glycosylated 19 kDa N-terminal fragment (Shh-N) and a glycosylated 25 kDa C-terminal protein (Shh-C) (4). Shh-C, which is responsible for the intramolecular processing of Shh, is rapidly degraded following Shh proteolysis (5). Shh-N is highly conserved, sharing >98% aa identity between mouse, human, rat, canine, porcine, and chicken Shh-N. Shh-N can be palmitoylated at its N-terminal cysteine and modified by cholesterol addition at its C-terminus (6). These modifications contribute to the membrane tethering of Shh as well as its assembly into various sized multimers (6-9). Lipid modification and multimerization greatly increase Shh-N receptor binding affinity and signaling potency (5, 6, 8, 9). Monomeric and multimeric Shh can be released from the plasma membrane by the cooperative action of DISP1, SCUBE2, and TACE/ADAM17 (10-12). Modifications also extend the effective range of Shh functionality and are required for the development of protein gradients important in tissue morphogenesis (9, 13). Canonical signaling of Shh is mediated by a multicomponent receptor complex that includes Patched (PTCH1, PTCH2) and Smoothened (SMO) (14). The binding of Shh to PTCH releases the basal repression of SMO by PTCH. Shh activity can also be regulated through interactions with heparin, glypicans, and membrane-associated Hip (hedgehog interacting protein) (13, 15, 16).

References:

1. Briscoe, J. and P.P. Thérond (2013) *Mol. Cell. Biol.* **14**:416.
2. Aviles, E.C. *et al.* (2013) *Front. Cell. Neurosci.* **7**:86.
3. Xie, J. *et al.* (2013) *OncoTargets Ther.* **6**:1425.
4. Echelard, Y. *et al.* (1993) *Cell* **75**:1417.
5. Zeng, X. *et al.* (2001) *Nature* **411**:716.
6. Feng, J. *et al.* (2004) *Development* **131**:4357.
7. Goetz, J.A. *et al.* (2006) *J. Biol. Chem.* **281**:4087.
8. Pepinsky, R.B. *et al.* (1998) *J. Biol. Chem.* **273**:14037.
9. Chen, M.-H. *et al.* (2004) *Genes Dev.* **18**:641.
10. Etheridge, L.A. *et al.* (2010) *Development* **137**:133.
11. Jakobs, P. *et al.* (2014) *J. Cell Sci.* **127**:1726.
12. Dierker, T. *et al.* (2009) *J. Biol. Chem.* **284**:8013.
13. Lewis, P.M. *et al.* (2001) *Cell* **105**:599.
14. Carpenter, D. *et al.* (1998) *Proc. Natl. Acad. Sci. USA* **95**:13630.
15. Filmus, J. and M. Capurro (2014) *Matrix Biol.* **35**:248.
16. Chuang, P.-T. and A.P. McMahon (1999) *Nature* **397**:617.