

DESCRIPTION

Source *E. coli*-derived
Ala1-Lys127, with and without an N-terminal Met
Accession # P48750.1

N-terminal Sequence Analysis Met & Ala1

Predicted Molecular Mass 14.7 kDa

SPECIFICATIONS

Activity Measured in a cell proliferation assay using DA3 mouse myeloma cells.
The ED₅₀ for this effect is 5-30 pg/mL.

Endotoxin Level <0.10 EU per 1 µg of the protein by the LAL method.

Purity >97%, by SDS-PAGE under reducing conditions and visualized by silver stain.

Formulation Supplied as a 0.2 µm filtered solution in PBS. See Certificate of Analysis for details.

PREPARATION AND STORAGE

Shipping The product is shipped with dry ice or equivalent. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage **Use a manual defrost freezer and avoid repeated freeze-thaw cycles.**

- 12 months from date of receipt, -20 to -70 °C as supplied.
- 1 month, 2 to 8 °C under sterile conditions after opening.

BACKGROUND

GM-CSF was initially characterized as a factor that can support the *in vitro* colony formation of granulocyte-macrophage progenitors. It is also a growth factor for erythroid, megakaryocyte, and eosinophil progenitors. GM-CSF is produced by a number of different cell types (including T cells, B cells, macrophages, mast cells, endothelial cells, fibroblasts, and adipocytes) in response to cytokine or inflammatory stimuli. On mature hematopoietic cells, GM-CSF is a survival factor for and activates the effector functions of granulocytes, monocytes/macrophages, and eosinophils (1, 2). GM-CSF promotes a Th1 biased immune response, angiogenesis, allergic inflammation, and the development of autoimmunity (3 - 5). It shows clinical effectiveness in ameliorating chemotherapy-induced neutropenia, and GM-CSF transfected tumor cells are utilized as cancer vaccines (6, 7). The 22 kDa glycosylated GM-CSF, similar to IL-3 and IL-5, is a cytokine with a core of four bundled α -helices (8 - 10). Mature rat GM-CSF shares 56% - 69% amino acid sequence identity with canine, feline, human, mouse, and porcine GM-CSF. GM-CSF exerts its biological effects through a heterodimeric receptor complex composed of GM-CSF R α /CD116 and the signal transducing common β chain (CD131) which is also a component of the high-affinity receptors for IL-3 and IL-5 (11, 12). In addition, GM-CSF binds a naturally occurring soluble form of GM-CSF R α (13). Rat GM-CSF is active on mouse cells, although mouse GM-CSF is only weakly active on rat cells (14, 15).

References:

1. Martinez-Moczygomba, M. and D.P. Huston (2003) *J. Allergy Clin. Immunol.* **112**:653.
2. Barreda, D.R. *et al.* (2004) *Dev. Comp. Immunol.* **28**:509.
3. Eksioğlu, E.A. *et al.* (2007) *Exp. Hematol.* **35**:1163.
4. Cao, Y. (2007) *J. Clin. Invest.* **117**:2362.
5. Fleetwood, A.J. *et al.* (2005) *Crit. Rev. Immunol.* **25**:405.
6. Heuser, M. *et al.* (2007) *Semin. Hematol.* **44**:148.
7. Hege, K.M. *et al.* (2006) *Int. Rev. Immunol.* **25**:321.
8. Kaushansky, K. *et al.* (1992) *Biochemistry* **31**:1881.
9. Diederichs, K. *et al.* (1991) *Science* **254**:1779.
10. Smith, L.R. *et al.* (1994) *Immunogenetics* **39**:80.
11. Onetto-Pothier, N. *et al.* (1990) *Blood* **75**:59.
12. Hayashida, K. *et al.* (1990) *Proc. Natl. Acad. Sci.* **87**:9655.
13. Pelley, J.L. *et al.* (2007) *Exp. Hematol.* **35**:1483.
14. Oaks, M.K. *et al.* (1995) *J. Interferon Cytokine Res.* **15**:1095.
15. Vandenberghe, P. *et al.* (1990) *Lymphokine Res.* **9**:381.