

DESCRIPTION

Source *E. coli*-derived
Ser31-Lys183
Accession # Q9D6Z6

N-terminal Sequence Analysis Ser31

Predicted Molecular Mass 17.4 kDa

SPECIFICATIONS

SDS-PAGE 16 kDa, reducing conditions

Activity Measured by its ability to induce IL-6 secretion by NIH-3T3 mouse embryonic fibroblast cells. Towne, J.E. *et al.* (2004) J. Biol. Chem. 279:13677.
The ED₅₀ for this effect is 1-6 ng/mL.

Endotoxin Level <0.01 EU per 1 μ g of the protein by the LAL method.

Purity >95%, by SDS-PAGE under reducing conditions and visualized by silver stain.

Formulation Lyophilized from a 0.2 μ m filtered solution in MES, NaCl, TCEP, EDTA, CHAPS and PEG 8000 with BSA as a carrier protein. See Certificate of Analysis for details.

PREPARATION AND STORAGE

Reconstitution Reconstitute at 100 μ g/mL in 10 mM Tris-HCl, pH 8.0.


Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

- 12 months from date of receipt, -20 to -70 °C as supplied.
- 1 month, 2 to 8 °C under sterile conditions after reconstitution.
- 3 months, -20 to -70 °C under sterile conditions after reconstitution.


DATA

Bioactivity



Recombinant Mouse IL-36 β /IL-1F8 (aa 31-183) (Catalog # 7060-ML) induces IL-6 secretion in the NIH-3T3 mouse embryonic fibroblast cell line. The ED₅₀ for this effect is 1-6 ng/mL.

SDS-PAGE



1 μ g/lane of Recombinant Mouse IL-36 β /IL-1F8 (aa 31-183) was resolved with SDS-PAGE under reducing (R) conditions and visualized by silver staining, showing a single band at 16 kDa.

BACKGROUND

Mouse interleukin-36 beta [IL-36 β ; previously IL-1F8, FIL-1 η (eta) and IL-1H2] is a member of the IL-1 family of proteins that includes IL-1 β , IL-1 α , IL-1ra, IL-18, IL-36Ra/IL-1F5, IL-36 α /IL-1F6, IL-37/IL-1F7, IL-36 γ /IL-1F9 and IL-1F10 (1 - 6). All family members show a 12 β -stranded β -trefoil configuration, share up to 50% amino acid (aa) sequence identity, and are believed to have arisen from a common ancestral gene (3, 4). Although two alternatively spliced transcript variants for human IL-36 β /IL-1F8 have been described, to date, only one mouse IL-36 β /IL-1F8 isoform is known (3). Mouse IL-36 β /IL-1F8 is synthesized as a 183 amino acid (aa) protein that contains no signal sequence, no prosegment and no potential N-linked glycosylation site(s) (1, 2). Mouse IL-36 β /IL-1F8 shares 61% and 74% aa identity with human IL-36 β isoform 2 and rat IL-36 β , respectively. IL-36 β is agonistic, stimulating release of inflammatory mediators such as IL-6 and IL-8, and cytotoxic peptides such as beta-defensins 2 and 3 that aid in defense against microbial pathogens (7 - 10). The receptor for IL-36 proteins is IL-1 Rrp2, with IL-1 RAcP as a coreceptor (7, 9). Antagonism of IL-36 proteins by IL-36Ra, which also binds IL-1 Rrp2, has been shown by some investigators (5, 6). Skin keratinocytes express highest levels of IL-36 proteins and their receptors, followed by epithelia in the esophagus, trachea and bronchae (7 - 9). IL-36 β expression is increased in psoriatic skin and may play a role in pathogenesis of psoriasis (7, 8). IL-36 β is also expressed in resting and activated monocytes and B cells, synovial fibroblasts, neurons and glia, and is detectable in plasma and body fluids (1, 7, 9, 11). IL-36 β , along with IL-36 α and IL-36 γ , is up-regulated by IL-1 α and TNF- α in keratinocytes, and has been shown to activate NF- κ B and MAPK signaling pathways in an IL-1 Rrp2-dependent manner (7 - 9). Full-length recombinant IL-36 proteins appear less active than their endogenous counterparts, but trimming of the N-termini enhances their activity (9, 12).

References:

1. Smith, D.E. *et al.* (2000) J. Biol. Chem. **275**:1169.
2. Kumar, S. *et al.* (2000) J. Biol. Chem. **275**:10308.
3. Nicklin, M.J.H. *et al.* (2002) Genomics **79**:718.
4. Dunn, E. *et al.* (2001) Trends Immunol. **22**:533.
5. Dinarello, C. *et al.* (2010) Nat. Immunol. **11**:973.
6. Barksby, H.E. *et al.* (2007) Clin. Exp. Immunol. **149**:217.
7. Towne, J.E. *et al.* (2004) J. Biol. Chem. **279**:13677.
8. Johnston, A. *et al.* (2011) J. Immunol. **186**:2613.
9. Magne, D. *et al.* (2005) Arthritis Res. Ther. **8**:R80.
10. van Asseldonk, E.J.P. *et al.* (2010) Obesity **18**:2234.
11. Wang, P. *et al.* (2005) Cytokine **29**:245.
12. Blumberg, H. *et al.* (2010) J. Immunol. **185**:4354.