

DESCRIPTION

Source Chinese Hamster Ovary cell line, CHO-derived
Ala253-Gly366
Accession # Q07104

N-terminal Sequence Analysis Ala253

Predicted Molecular Mass 13 kDa

SPECIFICATIONS

SDS-PAGE 14-23 kDa, reducing conditions

Activity Measured by its ability to induce Smad2 phosphorylation in P19 mouse embryonal carcinoma cells. Mazerbourg, S. *et al.* (2004) Mol. Endocrinol. 18:653.
50-100 ng/mL of Recombinant Mouse GDF-3 can effectively induce Smad2 phosphorylation.

Endotoxin Level <0.10 EU per 1 µg of the protein by the LAL method.

Purity >90%, by SDS-PAGE visualized with Silver Staining and quantitative densitometry by Coomassie® Blue Staining.

Formulation Lyophilized from a 0.2 µm filtered solution in HCl with BSA as a carrier protein. See Certificate of Analysis for details.

PREPARATION AND STORAGE

Reconstitution Reconstitute at 100 µg/mL in 4 mM HCl


Shipping The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

- 12 months from date of receipt, -20 to -70 °C as supplied.
- 1 month, 2 to 8 °C under sterile conditions after reconstitution.
- 3 months, -20 to -70 °C under sterile conditions after reconstitution.


DATA

SDS-PAGE



1 µg/lane of Recombinant Mouse GDF-3 (Catalog # 9009-GD) was resolved with SDS-PAGE under reducing (R) and non-reducing (NR) conditions and visualized by silver staining, showing R bands at 16 and 17 kDa and NR bands at 16 and 18 kDa.

Bioactivity



Recombinant Mouse GDF-3 (Catalog # 9009-GD) Induces Smad2 phosphorylation in P19 mouse embryonic carcinoma cells. P19 cells were incubated with 50 and 100 ng/ml of Recombinant Mouse GDF-3 for 60 minutes. The cells were lysed and Western blots were performed with anti-phospho Smad2 and eIF4E which was served as a loading control.

BACKGROUND

GDF-3 (previously called Vgr-2) is a TGF- β superfamily member belonging to the growth/differentiation factor family (1, 2). GDF-3 is expressed in undifferentiated embryonic stem (ES) cells, white adipose tissue and the brain (2-4). The 366 amino acid (aa) mouse GDF-3 contains a 22 aa signal sequence, a 230 aa propeptide and a 114 aa mature protein that contains one potential N-glycosylation site. The mature region contains a cysteine-knot structure that is conserved throughout family members. However, it lacks the fourth cysteine which is responsible for the formation of an inter-molecular disulfide bond, so GDF-3 may exist as a non-covalent homodimer (2, 5). Mature mouse GDF-3 shares 90% and 83% aa sequence identity with rat and human GDF-3, respectively. Most of GDF-3 is present as the uncleaved prepro form (6). The uncleaved and the mature forms both appear to have activity, but that activity may differ (5-8). All forms can oppose BMPs. In ES cells, inhibition of BMP2 signaling by GDF-3 maintains pluripotency (5, 7). GDF3 also influences early cell fate decisions; for example, deletion of mouse GDF-3 produces defects in the anterior visceral endoderm of the pre-gastrulation embryo (6-8). GDF-3 cooperates with GDF-1 in embryogenesis, and the mature protein has nodal-like activity (8, 9). Although GDF family members signal through BMP receptors (ALK1, 2, 3 and 6), which activate Smads 1, 5 and 8, GDF-3 signaling through ALK4 and ALK7, which activate Smads 2 and 3, has also been reported (9, 10). In adipocytes, GDF-3 is induced by a high fat diet, promoting adipogenesis and obesity (3, 10, 11).

References:

1. Levine, A.J. and A.H. Brivanlou (2006) *Cell Cycle* **5**:1069.
2. McPherron, A.C. and S.-J. Lee (1993) *J. Biol. Chem.* **268**:3444. Mouse cloning
3. Wang, W. *et al.* (2004) *Biochem. Biophys. Res. Comm.* **321**:1024.
4. Hexige, S. *et al.* (2005) *Neurosci. Lett.* **389**:83.
5. Levine, A.J. *et al.* (2009) *Dev. Biol.* **325**:43.
6. Levine, A.J. and A.H. Brivanlou (2005) *Development* **133**:209.
7. Peerani, R. *et al.* (2007) *EMBO J.* **26**:4744.
8. Chen, C. *et al.* (2006) *Development* **133**:319.
9. Andersson, O. *et al.* (2007) *Dev. Biol.* **311**:500.
10. Andersson, O. *et al.* (2008) *Proc. Natl. Acad. Sci. USA* **105**:7252.
11. Shen, J.J. *et al.* (2009) *Mol. Endocrinol.* **23**:113.