

DESCRIPTION

Source	Human embryonic kidney cell, HEK293-derived		
	Cynomolgus Monkey PD-L2/B7-DC (Leu20-Pro219) Accession # XP_005581839	IEGRMD	Human IgG ₁ (Pro100-Lys330)
	N-terminus		C-terminus

N-terminal Sequence Analysis Leu20

Structure / Form Disulfide-linked homodimer

Predicted Molecular Mass 49 kDa

SPECIFICATIONS

SDS-PAGE 62-77 kDa, reducing conditions

Activity Measured by its binding ability in a functional ELISA. When Recombinant Human PD-1 Fc Chimera (Catalog # 1086-PD) is immobilized at 1 µg/mL, 100 µL/well, the concentration of Recombinant Cynomolgus Monkey PD-L2/B7-DC Fc Chimera that produces 50% of the optimal binding response is approximately 0.1-0.6 µg/mL.

Endotoxin Level <0.10 EU per 1 µg of the protein by the LAL method.

Purity >95%, by SDS-PAGE visualized with Silver Staining and quantitative densitometry by Coomassie® Blue Staining.

Formulation Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details.

PREPARATION AND STORAGE


Reconstitution Reconstitute at 100 µg/mL in PBS.

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

- 12 months from date of receipt, -20 to -70 °C as supplied.
- 1 month, 2 to 8 °C under sterile conditions after reconstitution.
- 3 months, -20 to -70 °C under sterile conditions after reconstitution.

DATA



BACKGROUND

Programmed Death Ligand 2 (PD-L2), also known as B7-DC and butyrophilin-like protein, is a transmembrane member of the B7 family of proteins that provide signals for regulating T-cell activation and tolerance (1). Within the ECD, cynomolgous PD-L2 shares 71% and 96% aa sequence identity with mouse and human PD-L2, respectively (2, 3). PD-L2 is expressed on dendritic cells, subsets of activated CD4⁺ and CD8⁺ T cells, and memory B cells that differentiate into plasma cells (3-5). At inflammatory sites such as rheumatoid arthritis, allergen exposure, and virus infection, PD-L2 is up-regulated on synoviocytes, infiltrating macrophages, dendritic cells, and airway epithelial cells (6-10). PD-L2, along with B7-H1/PD-L1, binds to T cell PD-1 where it promotes IFN- γ production and CD40 Ligand up-regulation while inhibiting IL-4 production (2, 3, 11, 12). In addition, PD-L2 binds to RGM-B on macrophages and alveolar epithelial cells, supporting respiratory immune tolerance (13). In asthma, PD-L2 suppresses IL-5 and IL-13 production, promotes IL-12 production by dendritic cells, and supports allergen-induced airway hyper-responsiveness and mucus production (8, 10).

References:

1. Ceeraz, S. *et al.* (2013) Trends Immunol. **34**:556.
2. Latchman, Y. *et al.* (2001) Nat. Immunol. **2**:261.
3. Tseng, S.-Y. *et al.* (2001) J. Exp. Med. **193**:839.
4. Messal, N. *et al.* (2011) Mol. Immunol. **48**:2214.
5. Zuccarino-Catania, G.V. *et al.* (2014) Nat. Immunol. **15**:631.
6. Guo, G. *et al.* (2012) Clin. Rheumatol. **31**:271.
7. Loke, P. and J.P. Allison (2003) Proc. Natl. Acad. Sci. USA **100**:5336.
8. Matsumoto, K. *et al.* (2004) J. Immunol. **172**:2530.
9. Stanciu, L.A. *et al.* (2006) J. Infec. Dis. **193**:404.
10. Lewkowich, I.P. *et al.* (2013) Mucosal Immun. **6**:728.
11. Ghiotto, M. *et al.* (2010) Int. Immunol. **22**:651.
12. Shin, T. *et al.* (2003) J. Exp. Med. **198**:31.
13. Xiao, Y. *et al.* (2014) J. Exp. Med. **211**:943.