

DESCRIPTION

Source *Spodoptera frugiperda*, Sf 9 (baculovirus)-derived
Ala279-Ser390
Accession # P01137
Produced in an animal component free process (ACFP).

N-terminal Sequence Analysis Ala279

Structure / Form Disulfide-linked homodimer

Predicted Molecular Mass 12.8 kDa (monomer)

SPECIFICATIONS

SDS-PAGE 11 kDa, reducing conditions

Activity Measured by its ability to inhibit the IL-4-dependent proliferation of HT-2 mouse T cells. Tsang, M. *et al.* (1995) Cytokine 7:389. The ED₅₀ for this effect is 0.04-0.2 ng/mL. The specific activity of recombinant human TGF-β1 is approximately 2.5 x 10⁴ U/μg, which is calibrated against human TGF-β1 Standard (NIBSC code: 89/514).

Endotoxin Level <0.10 EU per 1 μg of the protein by the LAL method.

Purity >95%, by SDS-PAGE visualized with Silver Staining and quantitative densitometry by Coomassie® Blue Staining.

Formulation Lyophilized from a 0.2 μm filtered solution in Acetonitrile and TFA. See Certificate of Analysis for details.

PREPARATION AND STORAGE

Reconstitution Reconstitute at 100 μg/mL in sterile 4 mM HCl.

Shipping The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

- 12 months from date of receipt, -20 to -70 °C as supplied.
- 1 month, 2 to 8 °C under sterile conditions after reconstitution.
- 3 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

TGF-β1 (transforming growth factor beta 1) is one of three closely related mammalian members of the large TGF-β superfamily that share a characteristic cystine knot structure (1-7). TGF-β1, -2 and -3 are highly pleiotropic cytokines that are proposed to act as cellular switches that regulate processes such as immune function, proliferation and epithelial-mesenchymal transition (1-4). Each TGF-β isoform has some non-redundant functions; for TGF-β1, mice with targeted deletion show defects in hematopoiesis and endothelial differentiation, and die of overwhelming inflammation (2). Human TGF-β1 cDNA encodes a 390 amino acid (aa) precursor that contains a 29 aa signal peptide and a 361 aa proprotein (8). A furin-like convertase processes the proprotein to generate an N-terminal 249 aa latency-associated peptide (LAP) and a C-terminal 112 aa mature TGF-β1 (8, 9). Disulfide-linked homodimers of LAP and TGF-β1 remain non-covalently associated after secretion, forming the small latent TGF-β1 complex (8-10). Covalent linkage of LAP to one of three latent TGF-β binding proteins (LTBPs) creates a large latent complex that may interact with the extracellular matrix (9, 10). TGF-β is activated from latency by pathways that include actions of the protease plasmin, matrix metalloproteinases, thrombospondin 1 and a subset of integrins (10). Mature human TGF-β1 shares 100% aa identity with pig, dog and cow TGF-β1, and 99% aa identity with mouse, rat and horse TGF-β1. It demonstrates cross-species activity (1). TGF-β1 signaling begins with high-affinity binding to a type II ser/thr kinase receptor termed TGF-β RII. This receptor then phosphorylates and activates a second ser/thr kinase receptor, TGF-β RI (also called activin receptor-like kinase (ALK) -5), or alternatively, ALK-1. This complex phosphorylates and activates Smad proteins that regulate transcription (3, 11, 12). Contributions of the accessory receptors betaglycan (also known as TGF-β RIII) and endoglin, or use of Smad-independent signaling pathways, allow for disparate actions observed in response to TGF-β in different contexts (11).

References:

1. Derynck, R. and K. Miyazono (2008) Cold Spring Harbor Laboratory Press, 29.
2. Dunker, N. and K. Krieglstein (2000) Eur. J. Biochem. 267:6982.
3. Wahl, S.M. (2006) Immunol. Rev. 213:213.
4. Chang, H. *et al.* (2002) Endocr. Rev. 23:787.
5. Lin, J.S. *et al.* (2006) Reproduction 132:179.
6. Hinck, A.P. *et al.* (1996) Biochemistry 35:8517.
7. Mittl, P.R.E. *et al.* (1996) Protein Sci. 5:1261.
8. Derynck, R. *et al.* (1985) Nature 316:701.
9. Miyazono, K. *et al.* (1988) J. Biol. Chem. 263:6407.
10. Oklu, R. and R. Hesketh (2000) Biochem. J. 352:601.
11. de Caestecker, M. *et al.* (2004) Cytokine Growth Factor Rev. 15:1.
12. Zuniga, J.E. *et al.* (2005) J. Mol. Biol. 354:1052.

MANUFACTURING SPECIFICATIONS

Animal Component-Free Process (ACFP) Manufacturing Conditions

R&D Systems Animal Component-Free Process (ACFP) recombinant proteins are expressed in an animal-free certified *Sf 9* insect cell line using dedicated animal-free raw materials and labware. Production and purification procedures use equipment and media that are confirmed animal-free but performed outside our dedicated animal-free laboratories. Every stage of the manufacturing process follows R&D Systems' stringent Standard Operating Procedures (SOPs). The certified *Sf 9* insect cell bank has undergone extensive testing to certify the lack of cytopathogens by screening for various viruses, Mycoplasma, and Spiroplasmas using both *in vitro* and *in vivo* testing methods. For *ex vivo* research or bioproduction, [additional documentation](#) can be provided.

[Please read our complete ACFP Statement](#)