

DESCRIPTION	
Species Reactivity	Human
Specificity	Detects human IL-1 α /IL-1F1 in direct ELISAs and Western blots. In direct ELISAs, approximately 30% cross-reactivity with recombinant mouse (rm) IL-1 α is observed and less than 5% cross-reactivity with recombinant porcine IL-1 α , recombinant rat IL-1 α , recombinant cotton rat IL-1 α , recombinant human IL-1 β and rmlIL-1 β is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human IL-1 α /IL-1F1 Ser113-Ala271 Accession # Q53QF9
Endotoxin Level	<0.10 EU per 1 μ g of the antibody by the LAL method.
Formulation	Lyophilized from a 0.2 μ m filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 μ m filtered solution in PBS.

APPLICATIONS		
Please Note: Optimal dilutions should be determined by each laboratory for each application. <i>General Protocols</i> are available in the <i>Technical Information</i> section on our website.		
	Recommended Concentration	Sample
Western Blot	0.1 μ g/mL	Recombinant Human IL-1 α /IL-1F1 (Catalog # 200-LA)
Immunocytochemistry	5-15 μ g/mL	See Below
Neutralization	Measured by its ability to neutralize IL-1 α /IL-1F1-induced proliferation in the D10.G4.1 mouse helper T cell line. Symons, J.A. <i>et al.</i> (1987) in <i>Lymphokines and Interferons, a Practical Approach</i> . Clemens, M.J. <i>et al.</i> (eds): IRL Press. 272. The Neutralization Dose (ND ₅₀) is typically 4-20 ng/mL in the presence of 50 μ g/mL Recombinant Human IL-1 α /IL-1F1 and 1.25 μ g/mL concanavalin A.	

DATA			
<p>Neutralization</p> <p>The graph shows two curves. The x-axis represents Recombinant Human IL-1α/IL-1F1 concentration in pg/mL on a log scale from 10⁻² to 10². The y-axis represents Mean RFU from 2000 to 4000. An orange line with open circles shows protein-induced proliferation, which increases from ~2200 at 10⁻² pg/mL to ~3800 at 10² pg/mL. A green line with solid circles shows neutralization by the antibody, which starts at ~3800 RFU at 10⁻² ng/mL and decreases to ~2200 RFU at 10² ng/mL.</p>	<p>Cell Proliferation Induced by IL-1α/IL-1F1 and Neutralization by Human IL-1α/IL-1F1 Antibody. Recombinant Human IL-1α/IL-1F1 (Catalog # 200-LA) stimulates proliferation in the D10.G4.1 mouse helper T cell line in a dose-dependent manner (orange line). Proliferation elicited by Recombinant Human IL-1α/IL-1F1 (50 μg/mL) is neutralized (green line) by increasing concentrations of Goat Anti-Human IL-1α/IL-1F1 Antigen Affinity-purified Polyclonal Antibody (Catalog # AF-200-NA). The ND₅₀ is typically 4-20 ng/mL in the presence of concanavalin A (1.25 μg/mL).</p>	<p>Immunocytochemistry</p> <p>The image shows a field of cells with red and blue staining. A scale bar indicates 100.0 μm.</p>	<p>IL-1α/IL-1F1 in Human PBMCs. IL-1α/IL-1F1 was detected in immersion fixed human peripheral blood mononuclear cells (PBMCs) treated with LPS using Goat Anti-Human IL-1α/IL-1F1 Antigen Affinity-purified Polyclonal Antibody (Catalog # AF-200-NA) at 10 μg/mL for 3 hours at room temperature. Cells were stained using the NorthernLights™ 557-conjugated Anti-Goat IgG Secondary Antibody (red; Catalog # NL001) and counterstained with DAPI (blue). View our protocol for Fluorescent ICC Staining of Non-adherent Cells.</p>

PREPARATION AND STORAGE	
Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Interleukin 1 (IL-1) is a name that designates two proteins, IL-1 α and IL-1 β , which are the products of distinct genes, but which show approximately 25% amino acid sequence identity and which recognize the same cell surface receptors. Although IL-1 production is generally considered to be a consequence of inflammation, recent evidence suggests that IL-1 is also temporarily upregulated during bone formation and the menstrual cycle and can be induced in response to nervous system stimulation. In response to classic stimuli produced by inflammatory agents, infections or microbial endotoxins, a dramatic increase in the production of IL-1 by macrophages and various other cells is seen. Cells in particular known to produce IL-1 include osteoblasts, monocytes, macrophages, keratinocytes, Kupffer cells, hepatocytes, thymic and salivary gland epithelium, Schwann cells, fibroblasts and glia (oligodendroglia, astrocytes and microglia).

IL-1 α and IL-1 β are both synthesized as 31 kDa precursors that are subsequently cleaved into proteins with molecular weights of approximately 17,000 Da. Neither precursor contains a typical hydrophobic signal peptide sequence and most of the precursor form of IL-1 α remains in the cytosol of cells, although there is evidence for a membrane-bound form of the precursor form of IL-1 α . The IL-1 α precursor reportedly shows full biological activity in the EL-4 assay. Among various species, the amino acid sequence of mature IL-1 α is conserved 60% to 70% and human IL-1 has been found to be biologically active on murine cell lines. Both forms of IL-1 bind to the same receptors, designated type I and type II. Evidence suggests that only the type I receptor is capable of signal transduction and that the type II receptor may function as a decoy, binding IL-1 and thus preventing binding of IL-1 to the type I receptor.