

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human IL-5 in direct ELISAs and Western blots. In Western blots, approximately 25% cross-reactivity with recombinant mouse IL-5 is observed. Neutralizes the biological activity of recombinant IL-5 and will also neutralize the biolog
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>S. frugiperda</i> insect ovarian cell line Sf 21-derived recombinant human IL-5 Ile20-Ser134 Accession # P05113
Conjugate	Alexa Fluor 647 Excitation Wavelength: 650 nm Emission Wavelength: 668 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Neutralization	Optimal dilution of this antibody should be experimentally determined.
Western Blot	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Interleukin 5 is a T cell-derived factor that promotes the proliferation, differentiation and activation of eosinophils. In mice, IL-5 has also been shown to be a growth and differentiation factor for B cells. Various names previously used to describe IL-5 include: T cell replacing factor (TRF), B cell growth factor II (BCGFII), B cell differentiation factor μ (BCDF μ), eosinophil differentiation factor (EDF) and eosinophil colony-stimulating factor (E_o-CSF). Biologically active IL-5 is a disulfide-linked homodimer. The cDNAs for murine and human IL-5 encode precursor proteins with signal peptides that are cleaved to form mature proteins containing 113 and 115 amino acid residues, respectively. Murine and human IL-5 are 70% identical in their amino acid sequences and show species cross-reactivity. The genes for human and mouse IL-5 have been mapped to chromosome 5 and chromosome 11, respectively; closely linked to the genes for IL-3, IL-4 and GM-CSF.

IL-5 exerts its activity on target cells by binding to specific cell surface receptors. The functional high-affinity receptor for human IL-5 has been shown to be composed of a low-affinity IL-5 binding α -subunit and a non-binding common β -subunit that is shared with the high-affinity receptors for GM-CSF and IL-3.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.