

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human FGF-5 in direct ELISAs and Western blots. In direct ELISAs, less than 1% cross-reactivity with recombinant human (rh) FGF-4, rhFGF-6, rhFGF-7, recombinant mouse FGF-8b, rhFGF-9, rhFGF acidic, and rhFGF basic is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human FGF-5 Glu23-Gly268 (Lys238Asn, Pro245Ser) Accession # Q8NF90
Conjugate	Alexa Fluor 647 Excitation Wavelength: 650 nm Emission Wavelength: 668 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Neutralization	Optimal dilution of this antibody should be experimentally determined.
Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

The FGF family is comprised of at least seven polypeptides that are potent regulators of cell proliferation, differentiation and function. All FGFs have two conserved cysteine residues and share 30-50% sequence homology at the amino acid level. FGF-5 was originally identified as a transforming gene by the NIH-3T3 focus formation assay using DNA derived from human tumors. FGF-5 cDNA encodes a 267 amino acid residue protein with a putative 22 amino acid residue signal peptide. The murine homologue of FGF-5 was cloned and found to be 84% homologous to the human protein at the amino acid sequence level. Human and murine FGF-5 exhibit cross species activity.

In vitro, recombinant human FGF-5 is a mitogen for Balb/3T3 fibroblasts and bovine heart endothelial cells. FGF-5 was also reported to be a major muscle-derived survival factor for cultured spinal motoneurons. *In vivo*, FGF-5 is suggested to play important roles in both embryology and neurobiology. Developmentally, FGF-5 mRNA is initially found in the embryoblast followed by the lateral somatic mesoderm, where it may play a role in angiogenesis, plus the myotomes cranial to the tail region, where it may delay terminal myoblast differentiation during cell migration. FGF-5 continues to impact muscle post-natally where it is believed to function as a target-derived neurotrophic factor of skeletal muscle. In the nervous system, FGF-5 has been most often identified in neurons associated with the limbic system, notably in neurons of the olfactory bulb and pyramidal cells of the hippocampus. Hippocampal FGF-5 is suggested to serve as a neurotrophic and differentiative factor for cholinergic and serotonergic neurons projecting to this region.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.