

**DESCRIPTION**

<b>Species Reactivity</b>	Human
<b>Specificity</b>	Detects human CCL2/JE/MCP-1 in direct ELISAs and Western blots.
<b>Source</b>	Polyclonal Goat IgG
<b>Purification</b>	Antigen Affinity-purified
<b>Immunogen</b>	<i>E. coli</i> -derived recombinant human CCL2/JE/MCP-1 Gln24-Thr99 Accession # P13500
<b>Conjugate</b>	Alexa Fluor 405 Excitation Wavelength: 405 nm Emission Wavelength: 421 nm
<b>Formulation</b>	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide

\*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

**APPLICATIONS**

<b>Please Note:</b> Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.	
<b>Neutralization</b>	Optimal dilution of this antibody should be experimentally determined.
<b>Western Blot</b>	Optimal dilution of this antibody should be experimentally determined.
<b>Immunohistochemistry</b>	Optimal dilution of this antibody should be experimentally determined.

**PREPARATION AND STORAGE**

<b>Shipping</b>	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
<b>Stability &amp; Storage</b>	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

**BACKGROUND**

CCL2, also known as monocyte chemotactic and activating factor (MCAF), was initially purified independently by two groups based on its ability to chemoattract monocytes. The human CCL2 cDNA encodes a 99 amino acid residue precursor protein with a 23 residue hydrophobic signal peptide that is cleaved to generate the 76 residue mature protein. Natural CCL2 is heterogeneous in size due to the addition of O-linked carbohydrates and sialic acid residues. In addition to fibroblasts; tumor cells, smooth muscle cells, endothelial cells, and mononuclear phagocytes can also produce CCL2 either constitutively or upon stimulation by various stimuli. CCL2 is a member of the β (C-C) subfamily of chemokines. The existence of MCP-2 and MCP-3 with 62% and 73% amino acid identity respectively, to CCL2 have been reported. Consistent with it being a member of the chemokine β family, CCL2 has been shown to chemoattract monocytes. In addition, it will also activate monocytes to be cytostatic for some human tumor cell lines; to increase cytosolic free calcium; to generate and release monocyte superoxide anions and to release monocyte lysosomal enzymes *in vitro*. CCL2 was reported to be capable of regulating adhesion molecule expression and cytokine production in human monocytes as well as chemoattracting, activating, and inducing histamine release from basophils. The biological roles played by CCL2 in a number of inflammatory and non-inflammatory disease states characterized by the accumulation of leukocytes at the site of the lesion, including atherosclerosis, delayed hypersensitivity reactions, etc., are being determined. CCL2 can bind to the C-C chemokine receptor-1 that also binds MIP-1α, RANTES and MIP-1β. A specific receptor for CCL2 has also been cloned from THP-1 and MonoMac 6 cells.

**PRODUCT SPECIFIC NOTICES**

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.