

DESCRIPTION


Species Reactivity	Human
Specificity	Detects human CNTF R α in direct ELISAs and Western blots. In Western blots, greater than 50% cross-reactivity with recombinant rat CNTF sR α is observed and less than 5% cross-reactivity with recombinant human (rh) IL-2 sR γ and rhIL-6 sR is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>S. frugiperda</i> insect ovarian cell line Sf21-derived recombinant human CNTF R α Gln23-Pro346 (predicted) Accession # P26992
Endotoxin Level	<0.10 EU per 1 μ g of the antibody by the LAL method.
Formulation	Lyophilized from a 0.2 μ m filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 μ m filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	0.1 μ g/mL	Recombinant Human CNTF R α (Catalog # 303-CR)
Neutralization		Measured by its ability to neutralize CNTF R α -enhanced proliferation in the TF-1 human erythroleukemic cell line. The Neutralization Dose (ND ₅₀) is typically 10-20 μ g/mL in the presence of 1 μ g/mL Recombinant Human CNTF R α and 20 ng/mL Recombinant Human CNTF.

DATA



PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> • 12 months from date of receipt, -20 to -70 °C as supplied. • 1 month, 2 to 8 °C under sterile conditions after reconstitution. • 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

The high-affinity CNTF receptor complex, which mediates the biological action of CNTF, contains three proteins: the ligand-binding α subunit (CNTF R α), and the two signal-transducing proteins LIF R β and gp130. Whereas LIF R β and gp130 are widely expressed in many cell types, the expression of CNTF R α is restricted to the central and peripheral nervous systems. cDNAs encoding human and rat CNTF R α share 94% amino acid (aa) sequence identity. Human CNTF R α cDNA encodes a 372 aa precursor protein with a 22 aa residue signal peptide and four potential glycosylation sites. CNTF R α differs from other cytokine receptors in that it lacks transmembrane and cytoplasmic domains and is anchored to cell membranes by a glycosylphosphatidylinositol (GPI) linkage. Similar to other GPI-linked proteins, soluble CNTF receptor α (CNTF sR α) can be released from the cell surface by phosphatidylinositol-specific phospholipase C. CNTF sR α can be released from skeletal muscle in response to peripheral nerve injury and high concentrations of CNTF sR α have also been detected in human cerebrospinal fluid. CNTF sR α binds CNTF in solution and the complex can act on cells that express only LIF R β and gp130 but not CNTF R α .