

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human CNTF Rα in direct ELISAs and Western blots.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>S. frugiperda</i> insect ovarian cell line Sf 21-derived recombinant human CNTF Rα Gln23-Pro346 (predicted) Accession # P26992
Conjugate	Alexa Fluor 750 Excitation Wavelength: 749 nm Emission Wavelength: 775 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

Neutralization	Optimal dilution of this antibody should be experimentally determined.
Western Blot	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

The high-affinity CNTF receptor complex, which mediates the biological action of CNTF, contains three proteins: the ligand-binding α subunit (CNTF Rα), and the two signal-transducing proteins LIF Rβ and gp130. Whereas LIF Rβ and gp130 are widely expressed in many cell types, the expression of CNTF Rα is restricted to the central and peripheral nervous systems. cDNAs encoding human and rat CNTF Rα share 94% amino acid (aa) sequence identity. Human CNTF Rα cDNA encodes a 372 aa precursor protein with a 22 aa residue signal peptide and four potential glycosylation sites. CNTF Rα differs from other cytokine receptors in that it lacks transmembrane and cytoplasmic domains and is anchored to cell membranes by a glycosylphosphatidylinositol (GPI) linkage. Similar to other GPI-linked proteins, soluble CNTF receptor α (CNTF sRα) can be released from the cell surface by phosphatidylinositol-specific phospholipase C. CNTF sRα can be released from skeletal muscle in response to peripheral nerve injury and high concentrations of CNTF sRα have also been detected in human cerebrospinal fluid. CNTF sRα binds CNTF in solution and the complex can act on cells that express only LIF Rβ and gp130 but not CNTF Rα.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.