

DESCRIPTION


Species Reactivity	Human
Specificity	Detects human CXCL12/SDF-1 in direct ELISAs and Western blots.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human CXCL12/SDF-1 α Lys22-Lys89 Accession # Q6ICW0
Endotoxin Level	<0.10 EU per 1 μ g of the antibody by the LAL method.
Formulation	Lyophilized from a 0.2 μ m filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 μ m filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	0.1 μ g/mL	Recombinant Human/Feline CXCL12/SDF-1 β aa 19-93 (Catalog # 2716-SD) Recombinant Human/Feline CXCL12/SDF-1 β aa 22-93 (Catalog # 351-FS) Recombinant Human/Feline/Rhesus Macaque CXCL12/SDF-1 α (Catalog # 350-NS)
Neutralization		Measured by its ability to neutralize CXCL12/SDF-1 α -induced chemotaxis in the BaF3 mouse pro-B cell line transfected with human CXCR4. The Neutralization Dose (ND ₅₀) is typically 1.5-6.0 μ g/mL in the presence of 2 ng/mL Recombinant Human/Feline/Rhesus Macaque CXCL12/SDF-1 α .

DATA



PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> 12 months from date of receipt, -20 to -70 °C as supplied. 1 month, 2 to 8 °C under sterile conditions after reconstitution. 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

CXCL12, also known as SDF-1 (stromal cell-derived factor 1) and PBSF (Pre-B cell growth stimulatory factor), is a member of the CXC chemokine superfamily. Two alternatively spliced mRNAs encode CXCL12 α (89 aa residues) and CXCL12 β (93 aa residues) that differ by the addition of 4 amino acids at the carboxy-terminus. Human and mouse CXCL12 are highly conserved. Mature human and mouse CXCL12 differ only by 1 amino acid.