



DESCRIPTION	
<b>Species Reactivity</b>	Human
<b>Specificity</b>	Detects human CD4 in direct ELISAs and Western blots. In direct ELISAs, less than 5% cross-reactivity with recombinant mouse CD4 and recombinant rat CD4 is observed.
<b>Source</b>	Polyclonal Goat IgG
<b>Purification</b>	Antigen Affinity-purified
<b>Immunogen</b>	<i>S. frugiperda</i> insect ovarian cell line Sf 21-derived recombinant human CD4 Lys26-Trp390 Accession # P01730
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

**APPLICATIONS**

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
<b>Western Blot</b>	0.1 µg/mL	Recombinant Human CD4 (Catalog # 514-CD)
<b>Immunocytochemistry</b>	5-15 µg/mL	See Below
<b>Immunohistochemistry</b>	5-15 µg/mL	Immersion fixed paraffin-embedded sections of human tonsil

**DATA**

<p><b>Immunocytochemistry</b></p>  <p><b>CD4 in Human T Cells.</b> CD4 was detected in immersion fixed human T cells using 2 µg/mL Goat Anti-Human CD4 Antigen Affinity-purified Polyclonal Antibody (Catalog # AF-379-NA) for 3 hours at room temperature. Cells were stained (red) and counterstained (green). View our protocol for <a href="#">Fluorescent ICC Staining of Cells on Coverslips</a>.</p>	<p><b>Immunocytochemistry</b></p>  <p><b>CD4 in Human PBMCs.</b> CD4 was detected in immersion fixed human peripheral blood mononuclear cells (PBMCs) using Goat Anti-Human CD4 Antigen Affinity-purified Polyclonal Antibody (Catalog # AF-379-NA) at 10 µg/mL for 3 hours at room temperature. Cells were stained using the NorthernLights™ 557-conjugated Anti-Goat IgG Secondary Antibody (yellow; Catalog # NL001) and counterstained with DAPI (blue). View our protocol for <a href="#">Fluorescent ICC Staining of Non-adherent Cells</a>.</p>
---	---

**PREPARATION AND STORAGE**

<b>Reconstitution</b>	Reconstitute at 0.2 mg/mL in sterile PBS.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>● 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>● 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>● 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

**BACKGROUND**

CD4 is an approximately 55 kDa type I membrane glycoprotein that is expressed predominantly on most thymocytes and a subset of mature T lymphocytes. In humans, CD4 is also expressed to a lesser extent on monocytes and macrophage related cells. Human CD4 cDNA encodes a 458 amino acid (aa) residue precursor protein with a 25 aa residue signal peptide, a 371 aa residue extracellular region containing four immunoglobulin homology domains, a 24 aa residue transmembrane domain and a 38 aa residue cytoplasmic domain. CD4 is a coreceptor required for T cell recognition of antigens that are presented by class II major histocompatibility complexes. CD4 has been shown to be a coreceptor of HIV entry and specifically binds gp120, the external envelope glycoprotein of HIV.

**References:**

1. Capon, D.I. *et al.* (1991) *Annu. Rev. Immunol.* **9**:649.