

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse Lymphotoxin β R/TNFRSF3 in Western blots.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0 derived recombinant mouse Lymphotoxin beta Receptor Ser28-Pro218 Accession # P50284
Conjugate	Alexa Fluor 350 Excitation Wavelength: 346 nm Emission Wavelength: 442 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Lymphotoxin beta receptor (LT β R), also known as TNF RIII and TNF R-related protein (TNF Rrp) is a member of the TNF receptor superfamily, designated TNFRSF3. Human LT β R cDNA encodes a 435 amino acid (aa) residue type I membrane protein with a putative 30 aa residue signal peptide, a 193 aa residue extracellular domain and a 171 aa residue cytoplasmic domain. The extracellular domain of LT β R contains four cysteine-rich motifs characteristic of the TNF receptor superfamily. The cytoplasmic region of LT β R shares little sequence similarity with other TNF receptor family members, suggesting that different signaling mechanisms may be used. LT β R is expressed in a variety of tissues including visceral and lymphoid tissues. LT β R is also expressed by cell lines of monocytic, epithelial, and fibroblastic origins but not by T and B lymphocytes. Human and mouse LT β R share 76% aa sequence homology. The TNF family ligands that have been shown to bind and activate LT β R include LIGHT (also a ligand for HVEM) and the heterotrimeric lymphotoxin LT α 1/ β 2 or LT α 2/ β 1. Depending on the cell type, activation of LT β R has been shown to induce NF κ B activation, chemokine production, growth arrest, and apoptosis. In vivo, LT β R has been shown to play a critical role in controlling cellular immune functions and lymphoid organogenesis.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.