

## DESCRIPTION





<b>Species Reactivity</b>	Human
<b>Specificity</b>	Detects human Cathepsin D in direct ELISAs and Western blots. In direct ELISAs, approximately 20% cross-reactivity with recombinant mouse (rm) Cathepsin D and less than 5% cross-reactivity with recombinant human (rh) Cathepsin A, rhCathepsin C, rhCathepsin H, rhCathepsin L, and rhCathepsin Z is observed. In Western blots, approximately 20% cross-reactivity with rmCathepsin D and less than 1% cross-reactivity with rhBACE-1 is observed.
<b>Source</b>	Polyclonal Goat IgG
<b>Purification</b>	Antigen Affinity-purified
<b>Immunogen</b>	Mouse myeloma cell line NS0-derived recombinant human Cathepsin D Leu21-Leu412 Accession # P07339
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Western Blot</b>	1 µg/mL	See Below
<b>Immunohistochemistry</b>	5-15 µg/mL	See Below
<b>Immunoprecipitation</b>	25 µg/mL	Conditioned cell culture medium spiked with Recombinant Human Cathepsin D (Catalog # 1014-AS), see our available <a href="#">Western blot detection antibodies</a>
<b>Simple Western</b>	50 µg/mL	See Below

## DATA



## PREPARATION AND STORAGE

<b>Reconstitution</b>	Reconstitute at 0.2 mg/mL in sterile PBS.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"><li>● 12 months from date of receipt, -20 to -70 °C as supplied.</li><li>● 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li><li>● 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li></ul>

## BACKGROUND

Cathepsin D is a lysosomal aspartic protease of the pepsin family (1). Human cathepsin D is synthesized as a precursor protein, consisting of a signal peptide (residues 1-18), a propeptide (residues 19-64), and a mature chain (residues 65-412) (2-4). The mature chain can be processed further to the light (residues 65-161) and heavy (residues 169-412) chains. It is expressed in most cells and overexpressed in breast cancer cells (5). It is a major enzyme in protein degradation in lysosomes, and also involved in the presentation of antigenic peptides. Mice deficient in this enzyme showed a progressive atrophy of the intestinal mucosa, a massive destruction of lymphoid organs, and a profound neuronal ceroid lipofucinosi, indicating that cathepsin D is essential for proteolysis of proteins regulating cell growth and tissue homeostasis (6). Cathepsin D secreted from human prostate carcinoma cells are responsible for the generation of angiostatin, a potent endogenous inhibitor of angiogenesis (6).

### References:

1. Conner *et al.* in *Handbook of Proteolytic Enzymes* Barrett (1998) Academic Press, San Diego, p. 828.
2. Faust, *et al.* (1985) Proc. Natl. Acad. Sci. USA **82**:4910.
3. Westley and May (1987) Nucl. Acid Res. **15**:3773.
4. Redecker, *et al.* (1991) DNA Cell Biol. **10**:423.
5. Rochefort, *et al.* (2000) Clin. Chim. Acta. **291**:157.
6. Tsukuba, *et al.* (2000) Mol. Cells **10**:601.