

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Thrombopoietin R/Tpo R in direct ELISAs and Western blots. In direct ELISAs and Western blots, less than 1% cross-reactivity with recombinant human Epo R is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant human Thrombopoietin R/Tpo R Gln26-Tyr423 Accession # P40238
Conjugate	Alexa Fluor 750 Excitation Wavelength: 749 nm Emission Wavelength: 775 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Neutralization	Optimal dilution of this antibody should be experimentally determined.
Western Blot	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Thrombopoietin receptor (Tpo R), also known as myeloproliferative leukemia protein (c-mpl), is a 95 kDa type I transmembrane protein that is a member of the type I cytokine receptor family within the hematopoietin/cytokine receptor superfamily (1-4). The 635 amino acid (aa) full-length human Tpo R contains a 25 aa signal sequence, a 466 aa extracellular domain with a ligand binding domain and two fibronectin type III domains, a transmembrane (TM) domain and a cytoplasmic domain. The extracellular domain of human Tpo R shares 78%, 76%, 81%, 82% and 80% aa identity with mouse, rat, bovine, canine and equine Tpo R, respectively. Humans produce three distinct mRNA species; a P-form, a K-form, and a truncated form (Mpl-tr) lacking a TM domain (3-7). The P-form encodes the full-length receptor. The Mpl-tr form, which is expressed in both human and mouse, is intracellular and targets the P-form for degradation (5, 6). The 579 aa K-form has an alternate cytoplasmic domain, but does not dimerize with, or inhibit, the P-form (7). Thrombopoietin (Tpo) is a key regulator of megakaryocytopoiesis, thrombopoiesis and hematopoietic stem cell self-renewal, as reflected by expression of the Tpo R on megakaryocytes, platelets and hematopoietic progenitors (2, 8). Receptor dimerization occurs upon Tpo binding and initiates signaling through the Ras/MAP and JAK/STAT pathways (1, 2). Internalization and degradation of Tpo following Tpo R binding serves to downregulate circulating Tpo (9). Tpo R expressed at low levels on endothelial cells does not appear to contribute to regulation of Tpo (10). Inactivating mutations of Tpo R cause thrombocytopenia, and absence of functional Tpo R is lethal in humans, but not mice. Other mutations, including an activating change in the TM domain, can cause thrombocytosis (11, 12).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.