

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse Betacellulin/BTC in direct ELISAs and Western blots. In direct ELISAs and Western blots, approximately 10% cross-reactivity with recombinant human BTC is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant mouse Betacellulin/BTC Asp32-Gln118 Accession # Q543J8
Conjugate	Alexa Fluor 647 Excitation Wavelength: 650 nm Emission Wavelength: 668 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Neutralization	Optimal dilution of this antibody should be experimentally determined.
Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Betacellulin (BTC) is a member of the EGF family of cytokines which includes EGF, TGF- α , amphiregulin, HB-EGF, epiregulin, tomoregulin and the neuregulins. All EGF family members are synthesized as type I transmembrane precursor proteins containing one or more EGF-like domains in their extracellular region (1). BTC, a heparin-binding protein, was originally isolated from the conditioned media of mouse pancreatic beta tumor cells as a 32 kDa glycoprotein (2). The mouse BTC cDNA encodes a 177 amino acid (aa) residue precursor with a 31 aa signal peptide, an 87 aa residue extracellular region containing one EGF-like domain, a 21 aa transmembrane domain and a 38 aa cytoplasmic domain. Soluble BTC is released from the transmembrane precursor by proteolytic processing (3). Mouse BTC shares 93% and 79% aa sequence identity with rat and human BTC, respectively (1). The mouse BTC gene is tightly linked to that of amphiregulin on mouse chromosome 5 (4). BTC is expressed in most tissues including kidney, uterus, liver and pancreas. It is also present in body fluids including serum, milk and colostrum (5). BTC promotes pancreatic beta-cell growth and differentiation (6) and is a potent mitogen for retinal pigment epithelial cells, vascular smooth muscle cells and fibroblasts (1). The effects of BTC is mediated by binding to ErbB1 and ErbB4 homodimers as well as ErbB heterodimers (1).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.