## **R**Dsystems a biotechne brand

# Human Phospho-p53 (S15) Antibody

Antigen Affinity-purified Polyclonal Rabbit IgG Catalog Number: AF1043

SCRIPTIO

DESCRIPTION		
Species Reactivity	Human	
Specificity Detects human, monkey, and rat p53 when phosphorylated at S15 and mouse p53 when phosphorylated at S18.		
Source	Polyclonal Rabbit IgG	
Purification	Antigen Affinity-purified	
Immunogen	Phosphopeptide containing human p53 S15 site	
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.	

### APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.			
	Recommended Concentration	Sample	
Western Blot	0.2 µg/mL	See Below	
Immunocytochemistry	5-15 μg/mL	See Below	
Immunohistochemistry	5-15 μg/mL	See Below	
Immunoprecipitation	1 μg/500 μg cell lysate	CEM human T-lymphoblastoid cell line exposed to UV-C and MCF-7 human breast cancer cell line treated with camptothecin (CPT), see our available Western blot detection antibodies.	

DATA



### Detection of Human Phospho-p53 (S15) by Western Blot. Western blot shows lysates of CEM human Tlymphoblastoid cell line untreated or exposed to 100 J/m<sup>2</sup> UV-C for the indicated time. PVDF membrane was probed with 0.2 µg/mL Human Phospho-p53 (S15) Antigen Affinity-purified Polyclonal Antibody (Catalog # AF1043) followed by HRPconjugated Anti-Rabbit IaG Secondary Antibody (Catalog # HAF008). A specific band for Phospho-p53 (S15) was detected at approximately 53 kDa (as indicated). The phosphospecificity of this antibody was supported by decreased labeling following treatment with 600 U $\lambda$ phosphatase (λ-PPase) for 1 hour. For additional reference, duplicate lysates were probed with 1:5000 dilution Human/Mouse/Rat p53 HRPconjugated Antigen Affinitypurified Polyclonal Antibody (lower panel, Catalog # HAF1355). This experiment was conducted under reducing conditions and using Immunoblot Buffer Group 1.

### Immunocytochemistry



Untreated

Phospho-p53 (S15) in HeLa Human Cell Line. p53 phosphorylated at S15 was detected in immersion fixed HeLa human cervical epithelial carcinoma cell line treated with UV light (left panel) or untreated (right panel) using Rabbit Anti-Human Phospho-p53 (S15) Antigen Affinity-purified Polyclonal Antibody (Catalog # AF1043) at 10 µg/mL for 3 hours at room temperature. Cells were stained using the NorthernLights™ 557-conjugated Anti-Rabbit IgG Secondary Antibody (red; Catalog # NL004) and counterstained with DAPI (blue). Specific staining was localized to nuclei. Cells were costained using Rat Anti-Tubulin (Catalog # NB600-506, Novus Biologicals) and NorthernLights™ 493-conjugated Anti-Rat IgG Secondary Antibody (green; Catalog # NL015). View our protocol for Fluorescent ICC Staining of Cells on Coverslips.

Rev. 12/6/2018 Page 1 of 2

biotechne

Global bio-techne.com info@bio-techne.com techsupport@bio-techne.com TEL +1 612 379 2956 USA TEL 800 343 7475 Canada TEL 855 668 8722 China TEL +86 (21) 52380373 Europe | Middle East | Africa TEL +44 (0)1235 529449



# Human Phospho-p53 (S15) Antibody

Antigen Affinity-purified Polyclonal Rabbit IgG Catalog Number: AF1043

# Immunohistochemistry

Phospho-p53 (S15) in Human Breast Cancer Tissue. p53 phosphorylated at S15 was detected in immersion fixed paraffin-embedded sections of human breast cancer tissue using Human Phospho-p53 (S15) Antigen Affinitypurified Polyclonal Antibody (Catalog # AF1043) at 15 µg/mL overnight at 4 °C. Tissue was stained using the Anti-Rabbit HRP-DAB Cell & Tissue Staining Kit (brown; Catalog # CTS005) and counterstained with hematoxylin (blue). View our protocol for Chromogenic IHC Staining of immersion fixed paraffin-embedded Tissue Sections.

### Immunohistochemistry



Phospho-p53 (S15) in Human Breast Cancer Tissue. p53 phosphorylated at S15 was detected in immersion fixed paraffinembedded sections of human breast cancer tissue using Human Phospho-p53 (S15) Antigen Affinitypurified Polyclonal Antibody (Catalog # AF1043) at 15 µg/mL overnight at 4 °C. Tissue was stained using the Anti-Rabbit HRP-DAB Cell & Tissue Staining Kit (brown; Catalog # CTS005) and counterstained with hematoxylin (blue). View our protocol for Chromogenic IHC Staining of immersion fixed paraffin-embedded Tissue Sections.

PREPARATION AND STORAGE		
Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.	
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C	
Stability & Storage	<ul> <li>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</li> <li>12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>	

### BACKGROUND

The p53 tumor suppressor protein is a multi-functional transcription factor that regulates proliferation, cell cycle checkpoints, and apoptosis. The importance of p53 is underscored by its mutation in over 50% of human cancers. Human p53 is a 393 amino acid (aa) nuclear/cytoplasmic protein that contains an N-terminal activation domain, a specific DNA binding domain, and a C-terminal domain that mediates tetramer formation and regulates DNA binding. P53 is phosphorylated on Ser15 in response to irradiation, and this phosphorylation increases recruitment of the coactivator CREB-binding protein/p300. The ATM or ATR kinases can phosphorylate p53 at serine 15 (S15), which leads to cell cycle arrest. Serine 15 phosphorylation leads to p53 stabilization and enhances transactivation of p53 target genes.

Rev. 12/6/2018 Page 2 of 2



Global bio-techne.com info@bio-techne.com techsupport@bio-techne.com TEL +1 612 379 2956 USA TEL 800 343 7475 Canada TEL 855 668 8722 China TEL +86 (21) 52380373 Europe | Middle East | Africa TEL +44 (0)1235 529449