

## DESCRIPTION


<b>Species Reactivity</b>	Cotton Rat
<b>Specificity</b>	Detects cotton rat CCL3/MIP-1 $\alpha$ in direct ELISAs and Western blots. In direct ELISAs, approximately 50% cross-reactivity with recombinant mouse MIP-1 $\alpha$ is observed, 15% cross-reactivity with recombinant human (rh) MIP-1 $\alpha$ is observed, less than 10% cross-reactivity with rhMIP-1 $\beta$ , rh6Ckine, and rhHCC-4 is observed.
<b>Source</b>	Polyclonal Goat IgG
<b>Purification</b>	Antigen Affinity-purified
<b>Immunogen</b>	<i>E. coli</i> -derived recombinant cotton rat CCL3/MIP-1 $\alpha$ Ala24-Ala92 Accession # AAL26704
<b>Endotoxin Level</b>	<0.10 EU per 1 $\mu$ g of the antibody by the LAL method.
<b>Formulation</b>	Lyophilized from a 0.2 $\mu$ m filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 $\mu$ m filtered solution in PBS.

## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

	Recommended Concentration	Sample
<b>Western Blot</b>	0.1 $\mu$ g/mL	Recombinant Cotton Rat CCL3/MIP-1 $\alpha$ (Catalog # 1061-RM)
<b>Neutralization</b>	Measured by its ability to neutralize CCL3/MIP-1 $\alpha$ -induced chemotaxis in the BaF3 mouse pro-B cell line transfected with human CCR5. The Neutralization Dose (ND <sub>50</sub> ) is typically 0.1-0.4 $\mu$ g/mL in the presence of 10 ng/mL Recombinant Cotton Rat CCL3/MIP-1 $\alpha$ .	

## DATA



## PREPARATION AND STORAGE

<b>Reconstitution</b>	Reconstitute at 0.2 mg/mL in sterile PBS.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

**BACKGROUND**

MIP-1 $\alpha$  is a  $\beta$  family (CC) chemokine and has been designated CCL3. MIP-1 $\alpha$  and MIP-1 $\beta$ , two closely related but distinct proteins, were originally purified from medium conditioned by a LPS-stimulated murine macrophage cell line. Cotton rat MIP-1 $\alpha$  cDNA encodes a 92 amino acid (aa) residue precursor protein with a 23 aa putative signal peptide. Mature cotton rat MIP-1 $\alpha$  shares approximately 70% amino acid identity with human MIP-1 $\alpha$ . MIP-1 $\alpha$  is expressed in a wide variety of cells, including lymphocytes, fibroblasts, and epithelial cells, as well as monocytes/macrophages.

MIP-1 $\alpha$  has been shown to play an important role in the recruitment of mononuclear cells. Additionally, MIP-1 $\alpha$  has been reported to have chemoattractant and adhesive effects on lymphocytes, preferentially promoting the chemotaxis of Th1 cells. MIP-1 $\alpha$  has also been shown to attract B cells, eosinophils, and dendritic cells. In addition, MIP-1 $\alpha$  augments cytolytic activity of NK cells (1). MIP-1 $\alpha$  has been identified as a stem cell inhibitory factor that can inhibit the proliferation of hematopoietic stem cells *in vitro* as well as *in vivo*. It has been demonstrated that MIP-1 $\alpha$  can bind the chemokine receptors CCR1 and CCR5 (2).

**References:**

1. Robertson, M. (2002) J. Leukoc. Biol. **71**:173.
2. Zlotnik, A. *et al.* (2000) Immunity **12**:121.