

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse Epiregulin in direct ELISAs and Western blots. In direct ELISAs, approximately 40% cross-reactivity with recombinant human Epiregulin is observed. In Western blots, recombinant mouse Proepiregulin and mature Epiregulin is detected
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant mouse Epiregulin Val56-Leu101 Accession # Q61521
Conjugate	Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Neutralization	Optimal dilution of this antibody should be experimentally determined.
Western Blot	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Epiregulin is a member of the EGF family of growth factors which includes, among others, epidermal growth factor (EGF), transforming growth factor (TGF)-alpha, amphiregulin (ARG), HB (heparin-binding)-EGF, betacellulin, and the various heregulins. They are all synthesized as transmembrane precursors and converted to soluble forms by proteolytic cleavage. Epiregulin was originally purified from the mouse fibroblast-derived tumor cell line NIH3T3/T7 (1). The mouse Epiregulin cDNA encodes for a transmembrane precursor of 162 amino acid in length, with the mature soluble form comprising residues 56-101 (2). The mode of action of Epiregulin is similar to other EGF family members in that it binds to and activates the tyrosine-kinase, ErbB-family receptors (ErbB1 through B4) (3). Although it stimulates phosphorylation of all four receptors, it appears to interact primarily with ErbB1 and ErbB4. Epiregulin has the broadest specificity of the EGF-like ligands but seems to preferentially activate heterodimeric receptor complexes (4). Epiregulin exhibits a variety of biological effects. It was originally shown to both inhibit growth of several epithelial tumor cells and stimulate growth of fibroblasts and other types of cells (1). Epiregulin expression is upregulated in a number of carcinoma cell lines. It has also been shown to be an autocrine growth factor in human epidermal keratinocytes (5). Epiregulin has also been shown to play a role in the early steps of pregnancy, regulating attachment of the blastocyst to the uterine epithelium during the implantation process (6).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.