

DESCRIPTION

Species Reactivity	Cotton Rat
Specificity	Detects cotton rat CXCL10/IP-10/CRG-2 in direct ELISAs and Western blots. In direct ELISAs, less than 1% cross-reactivity with recombinant mouse CXCL-16 is observed. In Western blots, less than 5% cross-reactivity with recombinant human CXCL10
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant cotton rat CXCL10/IP-10/CRG-2 Thr27-Val94 Accession # AAL16935
Conjugate	Alexa Fluor 594 Excitation Wavelength: 590 nm Emission Wavelength: 617 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

IP-10, also named mob-1 in rat and crg-2 in mouse, is a member of the non-ELR-motif containing CXC chemokine subfamily and has been designated CXCL10. It is induced in various cell types (including keratinocytes, endothelial cells, fibroblasts, smooth muscle cells, monocytes/macrophages, astrocytes, neutrophils, hepatocytes, T cells, epithelial cells and mesothelial cells) in response to stimulation by type I and/or type II interferon as well as by LPS. IP-10 has been shown to chemoattract activated CD4+ and CD8+ T cells, activated NK cells and smooth muscle cells. It also induces the adhesion of activated T cells to VCAM and ICAM. In addition, IP-10 has been reported to antagonize the angiogenic activity of FGF. IP-10 binds to and activates CXCR3, the chemokine receptor that is shared by MIG/CXCL9 and I-TAC/CXCL11, two additional non-ELR CXC chemokines that are induced by interferon. Cotton rat IP-10 cDNA encodes a 94 amino acid (aa) residue precursor protein with a putative 21 aa signal peptide and a 73 aa mature protein with one potential N-linked glycosylation site. Mature cotton rat IP-10 shares 66%, 64%, and 66% aa sequence identity with rat, mouse and human IP10, respectively (1-7).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.