

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse Epigen in direct ELISAs and Western blots.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant mouse Epigen Leu53-Ala103 Accession # Q924X1.1
Conjugate	Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Epigen is an epithelial mitogen that belongs to the EGF superfamily (1). Mouse Epigen cDNA encodes a 153 amino acid (aa) residue type I transmembrane precursor protein with a 18 aa putative signal peptide, a 92 aa extracellular domain, a 20 aa transmembrane domain, and a 22 aa cytoplasmic domain. Expression of mouse Epigen has been detected in multiple tissues including testis, liver and heart. Over the entire protein, mouse Epigen shares from approximately 23-33% sequence homology with other EGF family members. The functional internal peptide containing the EGF motif however is more than 40% homologous with that of TGF- α and epiregulin. A human cDNA clone encoding a mouse epigen-like protein has also been identified (2). Similarly to other EGF family proteins, a soluble form of mouse Epigen can most likely be generated from the membrane-bound form by proteolytic cleavage. Although recombinant mouse Epigen has been shown to be mitogenic for fibroblasts (3) and epithelial cells (1), compared to EGF and TGF- α , it is much less potent. The EGF receptor ErbB1 can be activated upon mouse Epigen activation. Nevertheless, involvements of additional ErbB receptors in mouse Epigen signaling have not been ruled out.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.