

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Pref-1/DLK-1/FA1 in direct ELISAs and Western blots.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant human Pref-1 long isoform Ala24-Pro297 with Arg248Pro and Lys295Ser substitutions Accession # AAA75364
Conjugate	Alexa Fluor 350 Excitation Wavelength: 346 nm Emission Wavelength: 442 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunocytochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Pref-1 (Preadipocyte factor 1; also DLK-1 and FA1) is a 58-65 kDa member of the Notch/Serrata/Delta family of proteins. It is expressed in prechondrocytes and preadipocytes and appears to block progenitor cell differentiation into mature cell lineages. Mature human Pref-1 is a 360 amino acid (aa) type I transmembrane N- and O-linked glycoprotein. It contains a 280 aa extracellular region (aa 24-303), a 24 aa transmembrane segment (aa 304-327), and a 56 aa cytoplasmic domain (aa 328-383). The extracellular region contains six EGF-like domains, and undergoes proteolytic cleavage to generate a bioactive 50 kDa fragment, plus three 25-31 kDa fragments that show no activity. There are multiple potential splice variants. One shows a deletion of aa 229-301, a second possesses a six aa substitution for aa 1-52, a third shows a deletion of aa 210-277, while a fourth contains a six aa substitution for aa # 207-383. Over aa 24-297, human Pref-1 shares 82% aa identity with mouse Pref-1.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.