

DESCRIPTION





Species Reactivity	Human/Mouse
Specificity	Detects human Serpin F1/PEDF in direct ELISAs. Detects human and mouse Serpin F1/PEDF in Western blots.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human Serpin F1/PEDF Asp44-Pro418 Accession # P36955
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	0.2-2 µg/mL	See Below
Immunohistochemistry	5-15 µg/mL	See Below
Simple Western	2 µg/mL	See Below

DATA



PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> • 12 months from date of receipt, -20 to -70 °C as supplied. • 1 month, 2 to 8 °C under sterile conditions after reconstitution. • 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Serpin F1 (serine proteinase inhibitor-clade F #1; also PEDF, PIG35 and EPC-1) is a secreted, 50 kDa glycoprotein member of the clade F-subfamily, serpin superfamily of protease inhibitors. It is expressed by diverse cell types such as retinal pigment and breast epithelium, fibroblasts, astrocytes and hepatocytes. Serpin F1 circulates in blood and binds to type I collagen plus heparan sulfate. Although it is a serpin, it belongs to a class of serpins that are nonprotease inhibiting, and instead is known to possess potent antiangiogenic and neurotrophic activity. Human Serpin F1 is 418 amino acids (aa) in length. It contains a 19 aa signal sequence plus a 399 aa mature region that shows an NLS (aa 146-149), a neuroprotective motif (aa 354-359) and an antiangiogenesis segment (aa 387-411). Phosphorylation occurs at multiple sites that affect bioactivity. Over aa 44-418, human Serpin F1 shares 87% aa identity with mouse Serpin F1.