

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human EG-VEGF/PK1 in direct ELISAs and Western blots. In direct ELISAs and Western blots, approximately 10% cross-reactivity with recombinant rat EG-VEGF is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant human EG-VEGF/PK1 Ala20-Phe105 Accession # P58294
Conjugate	Alexa Fluor 350 Excitation Wavelength: 346 nm Emission Wavelength: 442 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Endocrine gland-derived vascular endothelial growth factor (EG-VEGF), also called prokineticin 1 (PK1), is a member of the prokineticin family of secreted proteins that share a common structural motif containing ten conserved cysteine residues that form five pairs of disulfide bonds (1, 2). Members of this family include the mammalian EG-VEGF/PK1 and PK2, as well as the venom protein A (VPRA) from the venom of black mamba snake and the frog *Bombina variegata*, Bv8 (1). Human EG-VEGF precursor is a 105 amino acid (aa) residue protein with a 19 aa signal peptide that is cleaved to yield a 86 aa mature protein (1, 2). EG-VEGF is expressed in multiple tissues including the gastrointestinal (GI) tract and steroidogenic glands (testis, ovary, placenta and adrenal glands). EG-VEGF has been shown to potently stimulate the contraction of GI smooth muscle. In addition, EG-VEGF is a tissue-specific angiogenic factor that exhibits biological activities similar to that of VEGF on select cells. It induces the proliferation, migration, and fenestration in cultured endocrine gland-derived capillary endothelial cells. EG-VEGF binds to and activates two closely related G protein-coupled receptors, EG-VEGF/PK1-R1 and EG-VEGF/PK2-R2 (3, 4). Activation of the receptors leads to stimulation of phosphoinositide turnover and activation of p44/p42 MAP kinase signaling pathways.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.