

DESCRIPTION

Species Reactivity	Zebrafish
Specificity	Detects zebrafish VEGF in direct ELISAs and Western blots. In Western blots, less than 2% cross-reactivity with recombinant human VEGF ₁₆₅ , recombinant mouse VEGF ₁₆₄ , and recombinant rat VEGF ₁₆₄ (under non-reducing)
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>S. frugiperda</i> insect ovarian cell line Sf 21-derived recombinant zebrafish VEGF ₁₆₅ Ala24-Arg188 Accession # O73682
Conjugate	Alexa Fluor 488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Neutralization	Optimal dilution of this antibody should be experimentally determined.
Western Blot	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Vascular endothelial growth factor (VEGF), also known as vascular permeability factor (VPF) and VEGF-A, is a potent mediator of both angiogenesis and vasculogenesis in the fetus and adult. It is a member of the PDGF family that is characterized by the presence of eight conserved cysteine residues. In human, at least eight alternate splice isoforms of VEGF-A, ranging from 206 amino acids (aa) to 121 aa in length, are known. In zebrafish, two VEGF isoforms, a 165 aa and a 121 aa isoform, have been reported. Mature zebrafish VEGF₁₆₅ shares 64%, 62% and 62% aa sequence identity with frog, human, and mouse VEGF₁₆₅, respectively. There are two tyrosine kinase receptors for VEGF reported in mammals termed VEGF R1 and VEGF R2/FLK-1. One receptor has been identified in zebrafish (FLK-1), and this may actually represent the orthologue to the early common ancestor for mammalian VEGF R1 and R2. All receptors are type I transmembrane proteins that show seven immunoglobulin-like domains extracellularly and a split kinase domain intracellularly. In addition to the tyrosine kinase receptors, neuropilin-1 (NRP-1) has been reported to be a coreceptor for VEGF binding. It is proposed that the presence of NRP-1 lowers the concentration of VEGF necessary for activation of VEGF R2. NRP-1 has been reported in both zebrafish and human. VEGF regulates multiple biological functions in endothelial cells, including cell proliferation, migration and survival. These functions of VEGF are mediated partly through the induction of nitric oxide and prostacyclin, as well as up-regulation of metalloproteinases. Together with other vascular-specific growth factors such as the Angiopoietins, VEGF have separate but complementary roles in angiogenesis and vasculogenesis (1-7).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.