

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse Frizzled-2 in direct ELISAs and Western blots. In direct ELISAs, approximately 30% cross-reactivity with recombinant mouse (rm) Frizzled-1 is observed and less than 5% cross-reactivity with rmFrizzled-4, recombinant human Friz
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant mouse Frizzled-2 Gln29-Leu168 Accession # Q9JIP6
Conjugate	Alexa Fluor 647 Excitation Wavelength: 650 nm Emission Wavelength: 668 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Wnt signaling is involved in variety of developmental processes including cell fate determination, cell polarity, tissue patterning and control of cell proliferation. Members of the Frizzled family of proteins serve as receptors for the Wnt signaling pathway. The founding member of this family was identified in *Drosophila* based on its role in tissue polarity in the adult cuticle and named for the disorganized appearance of bristle hairs on the mutant. Ten mammalian Frizzled genes have been identified to date. The predicted structure of Frizzled proteins is similar among all family members, containing a divergent N-terminal signal peptide, a highly conserved extracellular cysteine-rich domain, a variable-length linker region, a seven-pass transmembrane domain, and a variable-length C-terminal tail. One of the most conserved regions of the Frizzled proteins is the extracellular cysteine-rich domain (CRD) which spans approximately 120 AA and contains 10 invariant cysteines (1). Mouse Frizzled-2 shows 100% amino acid identity to human and rat Frizzled-2 in the CRD region. Frizzled-2 expression is greater in embryonic than adult tissues, with heart, brain, lung, kidney and gut showing the highest levels (2). In addition, Frizzled-2 is expressed during migration and proliferation of neural crest cells that populate the heart and great arteries (3). However, Frizzled-2 may not be strictly involved in developmental processes, as its expression is also upregulated in myofibroblasts during tissue repair after myocardial infarction (4). Frizzled-2 is implicated in the Wnt/Ca²⁺ pathway, a mechanism by which Wnt-5a signaling results in calcium release from intracellular stores and activation of PKC and calmodulin-dependent protein kinase (5). Secreted frizzled related proteins (sFRPs), antagonists of the Wnt pathway, have been shown to interact with both Wnt ligands and Frizzled receptors (6).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.