

DESCRIPTION


Species Reactivity	Human
Specificity	Detects human Rad17 when phosphorylated at S635.
Source	Polyclonal Rabbit IgG
Purification	Antigen Affinity-purified
Immunogen	Phosphopeptide containing human Rad17 S635 site
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS and Sodium Azide with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	0.5 µg/mL	See Below

DATA



PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	<p>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</p> <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

The DNA damage response activates cell cycle checkpoints to allow time for DNA repair and ensure the fidelity of each cell cycle. One group of proteins that transduce the DNA damage signal is the Rad family. The hRad17 protein is phosphorylated at S635 in cells exposed to multiple types of genotoxic stress, including ionizing radiation (IR) and ultraviolet light (UV).