

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human IL-26 in direct ELISAs and Western blots. In these formats, less than 5% cross-reactivity with recombinant human IL-10 is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human IL-26 (R&D Systems, Catalog # 1375-IL) Lys22-Gln171 Accession # Q9NPH9
Conjugate	Alexa Fluor 532 Excitation Wavelength: 534 nm Emission Wavelength: 553 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

IL-26 was originally cloned from herpesvirus saimiri (HVS)-transformed T cells and named AK155 (1). It is a member of the IL-10 family of class II cytokines that signal via heterodimeric receptor complexes composed of two type I transmembrane receptor subunits (2). The human IL-26 gene has been mapped to chromosome 12q15. It encodes a 171 amino acid polypeptide with a 21 amino acid signal peptide. In addition to HVS-transformed T cells, IL-26 is also expressed in other virus transformed T cell lines, fresh peripheral mononuclear cells, activated NK cells and T cells. A mouse homologue of human IL-26 has not been identified. IL-26 binds with high-affinity to the heterodimeric complex consisting of the ligand-binding IL-20 R α and non ligand-binding IL-10 R β (3). Activation of the receptor complex results in rapid phosphorylation of STAT1 and STAT3. Although the IL-26 receptor complex is highly specific for IL-26 and is not activated by other class II cytokines, the individual subunits of the IL-26 receptor complex are components in receptor complexes for other class II cytokines (1). IL-20 R α can form dimers with IL-20 R β to function as signaling receptors for IL-19, IL-20, and IL-24. IL-10 R β can complex with IL-10 R α , IL-22 R, and IL-28 R α to transduce signals for IL-10, IL-22, and the three novel IFNs (IL-28A, IL-28B and IL-29), respectively. The physiological functions of IL-26 remain to be determined. IL-26 was reported to be a homodimer in solution (1).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.